

Palladium-Catalyzed Fluoroalkylation via C(sp³)–S Bond Cleavage of Vinylsulfonium Salts

Yuan He, Zilong Huang, Juan Ma, Fei Huang, Jie Lin, Hongmei Wang,* Bao-Hua Xu, Yong-Gui Zhou,* and Zhengkun Yu*



Cite This: *Org. Lett.* 2021, 23, 6110–6114



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: An interrupted Pummerer/palladium-catalyzed fluoroalkylation strategy was developed for alkenyl C–H fluoroalkylthiolation. Palladium-catalyzed ring-opening fluoroalkylation via aliphatic C–S bond cleavage of the vinylsulfonium salts efficiently afforded fluoroalkylthiolated alkene derivatives from readily available alkene substrates and CsF. The protocol features broad substrate scopes and good functional group tolerance under an air atmosphere. The practicability of the synthetic method was demonstrated by transforming the multisubstituted alkene products to diverse fluoroalkylthiolated *N*-heterocycles.




Fluorine has been known to exist in many important chemicals.¹ By incorporating a fluorine atom or a fluoroalkyl group at a specific position in the candidate molecules, it is possible to fine-tune the pharmacological and pharmacokinetic properties of the drug candidates. Among the documented fluoroalkyl groups, fluoroalkylthio groups have been paid considerable attention due to their tunable lipophilicity, binding affinity, metabolic stability, and specific electronic properties.² Biologically active fluoroalkylthio motifs can be used as the key structural units for the construction of pharmaceutical agents such as M2 muscarinic receptor agonist,³ 5-HT2c receptor,⁴ and the inhibitor of cartilage matrix degradation.⁵ In the early days, harsh fluorinating reagents were used to prepare fluoroalkylated compounds.⁶ Because these reagents are potentially explosive, highly moisture-sensitive, and toxic, continuous efforts have been made to develop simple and green fluorination methods.⁷ A few fluorination methods have been reported for directly functionalizing aliphatic C–H bonds.⁸ In this regard, benzylic C–H⁹ and those positioned α to a carbonyl group¹⁰ can be fluorinated under relatively mild conditions. Decarboxylative fluorination has also been applied to construct a C(sp³)–F bond under radical or transition-metal catalysis conditions.¹¹ Recently, the difunctionalization of alkenes by fluorinating reagents has been paid much attention to access fluoroalkylated compounds.¹² The ring-opening fluorination of carbocycles has opened another route for the same purpose.^{13–16} Arylsulfonium salts have recently been employed as useful coupling partners not only in light of their accessibility from simple arenes but also due to their versatility for many carbon–carbon and carbon–heteroatom bond formation reactions.^{17,18} Notably, aryl thianthrenium and dibenzothio-phenonium salts have been successfully used for the site-selective fluorination of arenes (Scheme 1a);¹⁹ however,

vinylsulfonium salts have been paid much less attention due to the multiple reactivities of alkenyl moieties compared with their aryl analogs.²⁰


During the ongoing investigation of polar alkenes, we became interested in the regio- and stereoselective con-

Scheme 1. Fluoroalkylation Strategies

(a) Fluorination via arylsulfonium salts¹⁹



(b) This work: Fluoroalkylthiolation via vinylsulfonium salts



Received: June 30, 2021

Published: July 20, 2021




struction of multifunctionalized alkenes. In an attempt to conduct the vinylic C–H fluorination of α -alkenoyl ketene *N,S*-acetal **1Aa** with *N*-fluorobenzenesulfonimide (NFSI),²¹ we found that the vinylic C–H bond could be formally difluorinated to form **1Ba** (73%) (Scheme 1b). The use of α -aroyl analogs of **1Aa** led to similar results (see the Supporting Information (SI) for details), but its di-(methylthio)-substituted *S,S*-acetal analog did not react under the same conditions. Thus we envisioned that vinylic C–H fluorination might be realized via vinylsulfonium salts in a fashion similar to that of arylsulfonium salts.¹⁹ Unexpectedly, the reaction of a vinylsulfonium salt with CsF under palladium catalysis did not form the desired vinylic C–F bond through the cleavage of the vinylic C–S bond; instead, it underwent a fluoroalkylation process by cleavage of an aliphatic C–S bond. It has been known that *S*-alkyl tetrahydro-1*H*-thiophen-1-ium salts and analogs can undergo ring-opening reactions with nucleophiles such as thiolates, azide, halogens, amines, and so on.²² Herein we disclose an interrupted Pummerer/palladium-catalyzed fluoroalkylation strategy for vinylic C–H fluoroalkylation via vinylsulfonium salts (Scheme 1b).

Initially, the reaction of 1-(1,1-bis(methylthio)-3-oxo-3-phenyl-prop-1-en-2-yl)tetrahydro-1*H*-thiophen-1-ium trifluoromethanesulfonate (**1a**) with CsF was conducted to optimize the reaction conditions for the formation of 2-((4-fluorobutyl)thio)-3,3-bis(methylthio)-1-phenylprop-2-en-1-one (**2a**). Vinylsulfonium salt **1a** was conveniently prepared from the readily available alkene by the interrupted Pummerer reaction.^{18–20} (See the SI for details.) The reaction conditions were optimized to **1a**/CsF 1:2 (molar ratio), 5 mol % Pd(OAc)₂ as the catalyst, 1,4-dioxane as the solvent, 100 °C, 12 h under an air atmosphere, giving the desired product **2a** in 85% isolated yield. The use of pyridine·HF and NFSI soluble in organic solvents could not result in **2a**. The phase-transfer catalysts TEBAC (triethyl benzyl ammonium chloride) and 18-crown-6 remarkably diminished the reaction efficiency in the absence of the palladium catalyst, suggesting that the reaction does not merely proceed via a fluoride-promoted nucleophilic ring-opening pathway and the palladium catalysis plays a crucial role (Table S1). It is noteworthy that alkenyl fluoride **2a'** was not detected in the reaction mixture by ¹⁹F NMR analysis.

Under the optimal conditions, the scope of vinylsulfonium salts **1** generated from di(alkylthio)-substituted alkenes, that is, ketene dithioacetals, was explored (Scheme 2). They could exhibit diverse reactivity to form the fluoroalkylation products of type **2** in good to excellent yields. An obvious steric effect was observed for *ortho*-methyl-substituted α -benzoyl vinylsulfonium salts (**1b**–**1d**), and their reaction with CsF afforded products **2b**–**2d** (56–75%). Somehow, the α -(4-methoxy)-benzoyl-substituted substrate did not react to form the desired product **2e**. Halogens (F, Cl, and Br) and a CF₃ group on the α -benzoyl moiety did not show an obvious substituent effect, and the reaction formed products **2f**–**2k** in 81–87% yields. α -(2-Naphthoyl) and α -heteroaroyl (2-furoyl or 2-thienoyl)-substituted vinylsulfonium salts also reacted well to produce **2l**–**2n** in 82–87% yields. Di(ethylthio)-substituted vinylsulfonium salt (**1o**) reacted with CsF less efficiently than its di(methylthio) analog **1a**, yielding **2o** in 75% yield. In a similar fashion, cyclic five-membered alkylthio-substituted vinylsulfonium salts **1p**–**1r**, **1t**, and **1u** reacted to give the corresponding products **2p**–**2r**, **2t**, and **2u** (57–70%), whereas the 4-F substituent on the α -benzoyl moiety of **1s** obviously

Scheme 2. Scope of Sulfonium Salts of Ketene Dithioacetals (1)^a




^aConditions: **1** (0.30 mmol), CsF (0.60 mmol), Pd(OAc)₂ (0.015 mmol), dioxane (2 mL), air, 100 °C, 12 h. Yields refer to isolated products.

reduced the yield of **2s** to 34%. Cycloalkylthio-substituted α -(2-aphthoyl) (**1v**) and α -acetyl (**1w**) vinylsulfonium salts also smoothly underwent the reaction with CsF, leading to **2v** and **2w** (68–72%). However, the six-membered cycloalkylthio-substituted α -benzoyl vinylsulfonium salt **1x** enabled the formation of only **2x** in 45% yield, exhibiting a lower reactivity than its five-membered cycloalkylthio-substituted analog **1a**.

It should be noted that the molecular structure of compound **2l** was confirmed by the X-ray single-crystal crystallographic determination. (See the SI for details.)

The substituent and size effects from the cycloalkylsulfonium ring in **1** were then explored (Scheme 3a–c). The reaction of the vinylsulfonium salts derived from 2-methyltetrahydro-1*H*-thiophene, that is, vinylsulfonium salts **1y** and **1z**, with CsF gave a mixture of two inseparable fluoroalkylation products **2y/2y'** (51%, 10:3) and **2z/2z'** (39%, 2:1) via different aliphatic C–S bond cleavages of the cycloalkylsulfonium ring, which reveals that the sterically hindered C(sp³)–S bond is much easier to cleave. The vinylsulfonium salt of thietane **1z1** was decomposed quickly under the standard conditions. To our delight, the vinylsulfonium salts of tetrahydro-2*H*-thiopyran **1z2**–**1z4** efficiently reacted with CsF to produce the target products **2z2**–**2z4** (57–78%) bearing a C₅-fluoroalkylthio functionality, whereas products **2a**–**2z** contain a C₄-fluoroalkylthio chain. Unfortunately, further extension of the fluoroalkylthio chain failed because the vinylsulfonium salts corresponding to large aliphatic cyclic sulfoxides C_nH_{2n}S=O (*n* ≥ 6) could not be successfully prepared by the known methods. It is noteworthy that dibenzothiophene sulfonium salts of ketene dithioacetals **1** were not successfully prepared either due to the increased steric interaction in the interrupted Pummerer reaction.


Scheme 3. Diversity of the Fluoroalkylation



The reactivity of open-chain vinylsulfonium salts of type 3 was comparatively investigated (Scheme 3d–f). Treatment of vinylsulfonium salt **3a** derived from α -benzoyl ketene di(methylthio)acetal (**1A**) and dimethylsulfoxide (DMSO) with CsF under the standard conditions gave methylthiolated alkene **4a** in 91% isolated yield with methyl fluoride formed as the major byproduct. The parent alkene **1A** was generated in 6% yield from the decomposition of the starting vinylsulfonium salt, which might be promoted by moisture in air or by water present in the sulfonium salt during its preparation. In the case of the vinylsulfonium salt of methyl decyl thioether (**3b**), two kinds of C(sp³)–S bond cleavages occurred to form **4a** (16%) and **4b** (34%) by cleavage of the decyl–S and methyl–S bonds, respectively, as well as **1A** (31%). Notably, decyl fluoride was not detected in the reaction mixture by ¹⁹F NMR analysis, but MeF and methyl decyl thioether (RSMes) in both the gas and liquid phases were detected by GC–MS analysis. (See the SI for details.) Decyl triflate was presumably formed during the reaction because the corresponding molecular ion peak of decanal generated from the triflate byproduct was detected by GC–MS analysis. Compound **3c** behaved in a fashion similar to that of **3b**. Interestingly, the vinylsulfonium salt of methyl phenyl thioether (**3d**) predominantly underwent the decomposition reaction to form **1A** (52%) and **4a** (41%) with the release of PhSMe and PhOTf without phenyl fluoride formed in the reaction mixture. These results have suggested that the Me–S bond is easier to cleave than decyl/dodecyl–S

bonds in **3b/3c**, no fluoroalkylation occurred through the cleavage of these long-chain alkyl–S bonds, and the Ph–S bond is much easier to cleave than the Me–S bond in **3d** without the occurrence of fluoroarylation. In the case of using other counteranions such as BF₄[−], the chemoselectivities were not obviously affected, but the yields of **4a–4c** and **1A** were slightly decreased.

Next, the protocol generality was explored by performing the reaction of styryl sulfonium salts **5** with CsF (Scheme 4). In

Scheme 4. Scope of Styryl Sulfonium Salts (**5**)^a


^aConditions: **5** (0.30 mmol), CsF (0.60 mmol), Pd(OAc)₂ (0.015 mmol), dioxane (2 mL), air, 100 °C, 12 h. ^bPd(OAc)₂ (0.03 mmol), 24 h. ^cIsomer ratio was determined by ¹H NMR analysis. Yields refer to isolated products.

contrast with the multisubstituted vinylsulfonium salts of type 1, the sulfonium salt of styrene, that is, (*E*)-1-styryl-tetrahydro-1*H*-thiophen-1-ium trifluoromethanesulfonate (**5a**), reacted with CsF less efficiently, giving the target product **6a** in 12% yield. The yield was enhanced to 35% by increasing the catalyst loading to 10 mol % and extending the reaction time to 24 h. 1,1-Diaryl-substituted vinylsulfonium salts **5b–5h** underwent the reaction smoothly, efficiently yielding products **6b–6h** (61–79%). In the cases of vinylsulfonium salts **5c** and **5e** derived from 4-Me- and 3,4-Me₂-substituted 1,1-diaryl alkenes, isomeric products **6c** (61%, *E/Z* = 1:1) and **6e** (72%, *E/Z* = 9:5) were obtained, respectively. The use of 2-bromo-substituted 1,1-diaryl vinylsulfonium salt (*E*)-(**5d**) led to (*E*)-**6d** (64%), showing only one configuration. It is noteworthy that the sulfonium salt of 2-methylstyrene (**5i**) did not undergo the same type of fluoroalkylation reaction. The sulfonium salts of 1,2-stilbenes also reacted well with CsF, resulting in **6j** and **6k** (45–47%), exhibiting an obvious negative steric impact on the reaction efficiency. The reactivity difference is rationalized as follows. Styrylsulfonium salt **5a** was readily decomposed under the standard conditions, which led to **6a** in a low yield. An additional aryl at the one-position gives extra stabilization to 1,1-diarylvinylsulfonium salts **5b–5h** due to the conjugation effect, which reacted with CsF to afford the target products in decent yields. However, a negative steric impact exists in the sulfonium salts of 1,2-stilbenes (**5j** and **5k**), deteriorating the reaction efficiency. Interestingly, the 1,1-

diphenyl vinylsulfonium salt (**5I**) of tetrahydro-2*H*-thiopyran underwent ring-opening fluoroalkylation to afford the desired product **6I** (55%) bearing a C₅-fluoroalkylthio chain.

The gram-scale preparation of compound **2a** was performed on a 5 mmol scale of **1a** to give **2a** in 79% yield (1.3 g). In the presence of the K₂CO₃ base, the condensation of **2a** with guanidine nitrate formed 5-((4-fluorobutyl)thio)-4-(methylthio)-6-phenylpyrimidin-2-amine (**7**) in 75% yield (Scheme 5a). The treatment of **2a** with an excess of hydrazine hydrate

Scheme 5. Gram-Scale Experiment and Product Derivatization^a



^aReagents and conditions: (a) guanidine nitrate (2 equiv), K₂CO₃ (2 equiv), MeCN (2 mL), air, 100 °C, 36 h. (b) Hydrazine hydrate (10 equiv), EtOH (2 mL), air, 80 °C, 12 h. (c) Phenylhydrazine (2 equiv), EtOH (2 mL), air, 80 °C, 24 h. (d) Hydroxylamine hydrochloride (10 equiv), K₂CO₃ (10 equiv), EtOH (2 mL), air, 80 °C, 12 h.

or phenyl-hydrazine in refluxing ethanol afforded 4-fluoroalkylthiopyrazoles **8** and **9** in 90 and 81% yields, respectively (Scheme 5b,c). 4-((4-Fluorobutyl)thio)-3-(methylthio)-5-phenylisoxazole (**10**) was also efficiently synthesized (84%) by the reaction of **2a** and hydroxylamine (Scheme 5d). The molecular structures of compounds **7**–**9** were further confirmed by the X-ray single-crystal crystallographic determinations. (See the SI for details.)

Control experiments were conducted to probe into the reaction mechanism. The reaction of **1a** with CsF was carried out in the presence of 3.0 equiv of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) or 2,6-di-*tert*-butyl-4-methylphenyl (BHT) under the standard conditions. These radical-trapping reagents could not inhibit the ring-opening fluoroalkylation reaction, leading to **2a** in 37–55% yield, which excludes a radical pathway in the catalytic cycle. A palladium-catalyzed fluoroalkylation mechanism via aliphatic the C–S bond cleavage of vinylsulfonium salts is proposed. Because a small amount of the target product **2a** (11–15%) could be formed in the absence of the catalyst (Table S1), nucleophilic ring-opening fluorination cannot be excluded, but palladium-catalyzed ring-opening fluorination is predominant.

In conclusion, we have developed a C–H fluoroalkylthiolation strategy of alkenes via the palladium-catalyzed fluoroalkylation of vinylsulfonium salts under an air atmosphere. Multisubstituted fluoroalkylthiolated alkenes can be accessed by means of the ring-opening fluoroalkylation of the vinylsulfonium salts. The synthetic protocol has been shown to have potential for the synthesis of diverse fluoroalkylthiolated *N*-heterocycles.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.1c02172>.

Experimental materials and procedures, NMR of compounds, and X-ray crystallographic analysis for compounds **1Ba**, **2I**, and **7–9** (PDF)

Accession Codes

CCDC 2035272, 2073867, 2073869, 2077381, and 2091625 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Authors

Zhengkun Yu – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, P. R. China; Innovation Academy for Green Manufacture, Chinese Academy of Sciences, Beijing 100190, P. R. China; State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, P. R. China; orcid.org/0000-0002-9908-0017; Email: zkyu@dicp.ac.cn

Hongmei Wang – State Key Laboratory of NBC Protection for Civilian, Beijing 102205, P. R. China; orcid.org/0000-0003-1057-334X; Email: wanghongmei@dicp.ac.cn

Yong-Gui Zhou – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, P. R. China; orcid.org/0000-0002-3321-5521; Email: ygzhou@dicp.ac.cn

Authors

Yuan He – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, P. R. China; University of Chinese Academy of Sciences, Beijing 100049, P. R. China; orcid.org/0000-0001-6600-8433

Zilong Huang – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, P. R. China; University of Chinese Academy of Sciences, Beijing 100049, P. R. China; orcid.org/0000-0001-5350-8308

Juan Ma – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, P. R. China; University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Fei Huang – School of Food Science and Pharmaceutical Engineering, Nanjing Normal University, Nanjing 210023, P. R. China; orcid.org/0000-0002-5290-2490

Jie Lin – Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, P. R. China; University of Chinese Academy of Sciences, Beijing 100049, P. R. China

Bao-Hua Xu – Innovation Academy for Green Manufacture, Chinese Academy of Sciences, Beijing 100190, P. R. China; orcid.org/0000-0002-7222-4383

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.orglett.1c02172>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to the National Natural Science Foundation of China (21871253) for the support of our research.

■ REFERENCES

- (1) Fujiwara, T.; O'Hagan, D. Successful Fluorine-Containing Herbicide Agrochemicals. *J. Fluorine Chem.* **2014**, *167*, 16–29.
- (2) Wang, J.; Sánchez-Roselló, M.; Aceña, J.; del Pozo, C.; Sorochin-Sky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. Fluorine in Pharmaceutical Industry: Fluorine-Containing Drugs Introduced to the Market in the Last Decade (2001–2011). *Chem. Rev.* **2014**, *114*, 2432–2506.
- (3) Ma, Y.; Kiesewetter, D. O.; Jagoda, E. M.; Huang, B. X.; Eckelman, W. C. Identification of Metabolites of Fluorine-18-labeled M2Muscarinic Receptor Agonist, 3-(3-[(3-Fluoropropyl)thio]-1,2,5-thiadiazol-4-yl)-1,2,5,6-tetrahydro-1-methylpyridine, Produced by Human and Rat Hepatocytes. *J. Chromatogr. B: Anal. Technol. Biomed. Life Sci.* **2002**, *766*, 319–329.
- (4) Ladouceur, G. H.; Velthuisen, E.; Choi, S.; Zhang, Z.; Wang, Y.; Baryza, J. L.; Coish, P.; Smith, R.; Chen, J. Preparation of 4-Sulfanyl/Sulfonyl/Sulfonyl-1H-pyrazolyl Compounds for Use in Diseases Associated with the 5-HT_{2c} Receptor. WO 2003057674A1, 2003.
- (5) Yasunori, F.; Masayuki, T.; Shinji, M.; Kohei, N. Preparation of Fused Imidazolidine Derivatives as Inhibitors of Cartilage Matrix Degradation. WO2002092606A1, 2002.
- (6) Kirk, K. L. Fluorination in Medicinal Chemistry: Methods, Strategies, and Recent Developments. *Org. Process Res. Dev.* **2008**, *12*, 305–321.
- (7) Campbell, M. G.; Ritter, T. Modern Carbon–Fluorine Bond Forming Reactions for Aryl Fluoride Synthesis. *Chem. Rev.* **2015**, *115*, 612–633.
- (8) Liu, W.; Huang, X.; Cheng, M.-J.; Nielsen, R. J.; Goddard, W. A., III; Groves, J. T. Oxidative Aliphatic C–H Fluorination with Fluoride Ion Catalyzed by a Manganese Porphyrin. *Science* **2012**, *337*, 1322–1325.
- (9) Lee, S. J.; Brooks, A. F.; Ichiishi, N.; Makaravage, K. J.; Mossine, A. V.; Sanford, M. S.; Scott, P. J. H. C–H ¹⁸F-Fluorination of 8-Methylquinolines with Ag[¹⁸F]F. *Chem. Commun.* **2019**, *55*, 2976–2979.
- (10) Kitamura, T.; Muta, K.; Muta, K. Hypervalent Iodine-Promoted α -Fluorination of Acetophenone Derivatives with a Triethylamine-HF Complex. *J. Org. Chem.* **2014**, *79*, 5842–5846.
- (11) Yin, F.; Wang, Z.; Li, Z.; Li, C. Silver-Catalyzed Decarboxylative Fluorination of Aliphatic Carboxylic Acids in Aqueous Solution. *J. Am. Chem. Soc.* **2012**, *134*, 10401–10404.
- (12) Chen, F.; Xu, X.-H.; Qing, F.-L. Photoredox-Catalyzed Addition of Dibromofluoromethane to Alkenes: Direct Synthesis of 1-Bromo-1-fluoroalkanes. *Org. Lett.* **2021**, *23*, 2364–2369.
- (13) Zhao, H.; Fan, X.; Yu, J.; Zhu, C. Silver-Catalyzed Ring-Opening Strategy for the Synthesis of β - and γ -Fluorinated Ketones. *J. Am. Chem. Soc.* **2015**, *137*, 3490–3493.
- (14) Ishida, N.; Okumura, S.; Nakanishi, Y.; Murakami, M. Ring-opening Fluorination of Cyclobutanols and Cyclopropanols Catalyzed by Silver. *Chem. Lett.* **2015**, *44*, 821–823.
- (15) Pitts, C. R.; Ling, B.; Snyder, J. A.; Bragg, A. E.; Lectka, T. Aminofluorination of Cyclopropanes: A Multifold Approach through a Common, Catalytically Generated Intermediate. *J. Am. Chem. Soc.* **2016**, *138*, 6598–6609.
- (16) Wang, M.-M.; Waser, J. Oxidative Fluorination of Cyclopropylamides through Organic Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2020**, *59*, 16420–16424.
- (17) (a) Kaiser, D.; Klose, I.; Oost, R.; Neuhaus, J.; Maulide, N. Bond-Forming and -Breaking Reactions at Sulfur(IV): Sulfoxides, Sulfonium Salts, Sulfur Ylides, and Sulfinates. *Chem. Rev.* **2019**, *119*, 8701–8780. (b) Yorimitsu, H. Cascades of Interrupted Pummerer Reaction-Sigmatropic Rearrangement. *Chem. Rec.* **2017**, *17*, 1156–1167. (c) Nogi, K.; Yorimitsu, H. Aromatic Metamorphosis: Conversion of An Aromatic Skeleton into A Different Ring System. *Chem. Commun.* **2017**, *53*, 4055–4065.
- (18) (a) Cowper, P.; Jin, Y.; Turton, M. D.; Kociok-Köhn, G.; Lewis, S. E. Azulen sulfonium Salts: Accessible, Stable, and Versatile Reagents for Cross-Coupling. *Angew. Chem., Int. Ed.* **2016**, *55*, 2564–2568. (b) Berger, F.; Plutschack, M. B.; Riegger, J.; Yu, W. W.; Speicher, S.; Ho, M.; Frank, N.; Ritter, T. Site-selective and Versatile Aromatic C–H Functionalization by Thianthrenation. *Nature* **2019**, *567*, 223–228. (c) Aukland, M. H.; Siauciulis, M.; West, A.; Perry, G. J. P.; Procter, D. J. Metal-free Photoredox-catalysed Formal C–H/C–H Coupling of Arenes Enabled by Interrupted Pummerer Activation. *Nat. Catal.* **2020**, *3*, 163–169. (d) Wang, S.-M.; Song, H.-X.; Wang, X.-Y.; Liu, N.; Qin, H.-L.; Zhang, C.-P. Palladium-Catalyzed Mizoroki–Heck-Type Reactions of [Ph₂SR_{fn}][OTf] with Alkenes at Room Temperature. *Chem. Commun.* **2016**, *52*, 11893–11896. (e) Tian, Z.-Y.; Zhang, C.-P. Ullmann-Type N-Arylation of Anilines with Alkyl(aryl)sulfonium Salts. *Chem. Commun.* **2019**, *55*, 11936–11939.
- (19) Li, J. K.; Chen, J.; Sang, R.; Ham, W.-S.; Plutschack, M. B.; Berger, F.; Chhabra, S.; Schnegg, A.; Genicot, C.; Ritter, T. Photoredox Catalysis with Aryl Sulfonium Salts Enables Site-Selective Late-Stage Fluorination. *Nat. Chem.* **2020**, *12*, 56–62.
- (20) (a) Aukland, M. H.; Talbot, F. J. T.; Fernandez-Salas, J. A.; Ball, M.; Pulis, A. P.; Procter, D. J. An Interrupted Pummerer/Nickel-Catalysed Cross-Coupling Sequence. *Angew. Chem., Int. Ed.* **2018**, *57*, 9785–9789. (b) Chen, J. T.; Li, J. K.; Plutschack, M. B.; Berger, F.; Ritter, T. Regio- and Stereoselective Thianthrenation of Olefins To Access Versatile Alkenyl Electrophiles. *Angew. Chem., Int. Ed.* **2020**, *59*, 5616–5620. (c) He, Z.; Shrivs, H. J.; Fernández-Salas, J. A.; Abengózar, A.; Neufeld, J.; Yang, K.; Pulis, A. P.; Procter, D. J. Synthesis of C2 Substituted Benzothiophenes via an Interrupted Pummerer/[3,3]-Sigmatropic/1,2-Migration Cascade of Benzothiophene S–Oxides. *Angew. Chem., Int. Ed.* **2018**, *57*, 5759–5764. (d) Zhang, Y.-L.; Yang, L.; Wu, J.; Zhu, C. Y.; Wang, P. Vinyl Sulfonium Salts as the Radical Acceptor for Metal-Free Decarboxylative Alkenylation. *Org. Lett.* **2020**, *22*, 7768–7772.
- (21) Zupan, M.; Iskra, J.; Stavber, S. Chemistry of Organo Halogenic Molecules. 140. Role of the Reagent Structure on the Transformations of Hydroxy Substituted Organic Molecules with the N-Fluoro Class of Fluorinating Reagents. *Bull. Chem. Soc. Jpn.* **1995**, *68*, 1655–1660.
- (22) (a) Eliel, E. L.; Hutchins, R. O.; Mebane, R.; Willer, R. L. Endocyclic vs. Exocyclic Attack in Nucleophilic Displacement Reactions on Five- and Six-Membered Cyclic Onium Salts. *J. Org. Chem.* **1976**, *41*, 1052–1057. (b) Benn, M. H.; Singh, V. K. A Simple, Biogenetically Modeled Synthesis of 4-(Methylthio)butylthiocyanate: The Reaction of Thiocyanate Anion with S-Methyl-(1, n)-epithioniums. *Can. J. Chem.* **1986**, *64*, 940–942. (c) Vo, D.-V.; Truong, V.-D.; Tran, T.-D.; Do, V.-T.-N.; Pham, N.-T.-A.; Thai, K.-M. A New and Effective Approach to the Synthesis of Sulfuraphane. *Lett. Org. Chem.* **2015**, *13*, 7–10. (d) Wang, N.; Jia, Y. M.; Qin, H. M.; Jiang, Z.-X.; Yang, Z. G. Chlorotrifluoromethyl-thiolation of Sulfur Ylides for the Formation of Tetrasubstituted Trifluoromethylthiolated Alkenes. *Org. Lett.* **2020**, *22*, 7378–7382. (e) Lopez-Alled, C. M.; Martin, F. J. O.; Chen, K.-Y.; Kociok-Köhn, G.; James, T. D.; Wenk, J.; Lewis, S. E. Azulen sulfonium and Azulen bis(sulfonium) Salts: Formation by Interrupted Pummerer Reaction and Subsequent Derivatization by Nucleophiles. *Tetrahedron* **2020**, *76*, 131700–131711.