

Hypoiodite-Catalyzed Oxidative Umpolung of Indoles for Enantioselective Dearomatization

Hiroki Tanaka, Naoya Ukegawa, Muhammet Uyanik,* and Kazuaki Ishihara*



Cite This: *J. Am. Chem. Soc.* 2022, 144, 5756–5761



Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: Here we report the oxidative umpolung of 2,3-disubstituted indoles toward enantioselective dearomative aza-spirocyclization to give the corresponding spiroindolenines using chiral quaternary ammonium hypoiodite catalysis. Mechanistic studies revealed the umpolung reactivity of C3 of indoles by iodination of the indole nitrogen atom. Moreover, the introduction of pyrazole as an electron-withdrawing auxiliary group at C2 suppressed a competitive dissociative racemic pathway, and enantioselective spirocyclization proceeded to give not only spiropyrrolidines but also four-membered spiroazetidines that are otherwise difficult to access.


Indole-derived alkaloids are the largest group of nitrogen-containing secondary metabolites.¹ Because of their wide range of biological and pharmacological activities, a tremendous amount of research has been devoted to the development of efficient methods for the synthesis of indole alkaloids in synthetic organic chemistry, especially for drug discovery.^{1c,2} From this perspective, dearomatization of structurally planar indoles into enantioenriched three-dimensional indoline or indolenine structures has emerged as a powerful tool for the asymmetric synthesis of indole-derived alkaloids.³ Because of the inherent nucleophilicity of indoles, especially at C3 due to enamine-like reactivity, dearomatization reactions often proceed by addition of electrophiles at C3 (Scheme 1a, left).^{3c,d,4}

Conventionally, the incorporation of strong electron-withdrawing substituents^{5b,6} or leaving groups⁷ on the indole nucleus is often required for inversion of the polarity (umpolung)⁸ of nucleophilic indoles to give electrophilic

indoles. Recently, several asymmetric transformations of indole derivatives based on the C3 umpolung reactivity of 2-indolylmethanols have been achieved using chiral Brønsted acid catalysis.^{7b–e} On the other hand, methods for direct umpolung without preactivation would open a new avenue for the dearomative functionalization of indoles (Scheme 1a, right).^{5a} However, only a few examples of oxidative umpolung of indoles have been reported.^{9–11} Recently, new methods, namely electro¹⁰ and photochemical¹¹ oxidation of indoles, have been developed. Enantioselective dearomative coupling of indoles has also been achieved in combination with chiral phosphate or phosphoric acid catalysts under photooxidation conditions. However, these methods have relied on the use of transition metal chromophores as photocatalyst as well as preoxidized nucleophiles such as nitroxyl radicals or hydroxylamine derivatives.^{11a,b}

We recently reported the quaternary ammonium hypoiodite-catalyzed¹² oxidative C2-cyclization/peroxidation of homotryptamine derivatives (Scheme 1b).¹³ We envisioned that the introduction of a substituent at C2 of homotryptamines might sterically suppress cyclization or 1,2-migration¹⁴ at C2, allowing spirocyclization to proceed at C3 to form five-membered rings. Dearomative spirocyclization of homotryptamine derivatives to form aza-spiroindolenines has been reported.¹⁵ However, preinstallation of an O-based leaving group on an electrophilic nitrogen tether^{15a–c} or stoichiometric amounts of an organoiodine(III) as an oxidant^{15d} were required. In addition, enantioselective dearomative aza-spirocyclization remains elusive. Here we report the oxidative umpolung of 2,3-disubstituted indoles toward enantioselective

Scheme 1. Oxidative Umpolung of Indoles for Dearomative Coupling Reactions



Received: February 17, 2022


Published: March 23, 2022



dearomatizative spirocyclization of not only homotryptamines but also tryptamines to afford the corresponding spiroindolenines using quaternary ammonium hypoiodite catalysis^{12a,16} (Scheme 1c). Mechanistic studies revealed the umpolung reactivity of indole by iodination at N1.

We began our investigation by examining the oxidative spirocyclization of C2-methyl substituted homotryptamine derivative **1a** using *tert*-butyl hydroperoxide (TBHP) as an oxidant in the presence of 10 mol % *n*-tetrabutylammonium iodide (Bu₄NI) (Scheme 2a). Although the desired spiroindo-

Scheme 2. Investigation of Reaction Conditions and Chiral Catalysts




lene **2a** was obtained in 31% yield, intermolecular coupling with TBHP also proceeded to give peroxyindolenine **3** as a byproduct. To suppress this side reaction, we used cumene hydroperoxide (CHP) as a more sterically hindered oxidant instead of TBHP, and to our delight, a cleaner reaction was observed to give **2a** in 67% yield.¹⁷

Next, we investigated enantioselective spirocyclization using bis(binaphthyl)-based chiral quaternary ammonium iodide **4a**, (Scheme 2b).^{12a,16,18} After optimization of the reaction conditions (Table S1),¹⁹ **2a** was obtained in 70% yield with 46% ee in a methyl *tert*-butyl ether (MTBE)/toluene mixed solvent. Toluene and MTBE were found to be effective as solvents to enhance the reactivity and enantioselectivity, respectively.

Since investigation of the substituents at the 3- and 3'-positions of bis(binaphthyl)ammonium cation **4** failed to improve the enantioselectivity, we explored the use of mono(binaphthyl)-based catalysts **4**¹⁸ (Table S3). The use of morpholine-derived mono(binaphthyl)ammonium iodides improved the enantioselectivity, and the best result (81% ee) was obtained with *cis*-2,6-dimethylmorpholine-derived **4b** (Scheme 2b). To further improve the chemoselectivity, we used Na₂SO₄ as a desiccant to give **2a** in 87% yield with 81% ee (Table S4). The absolute stereochemistry of **2a** was assigned to be *R* by X-ray analysis of an enantiomerically pure sample that was obtained after a single recrystallization.

We examined the enantioselective dearomatative aza-spirocyclization of several *N*-(4-nosyl)homotryptamines **1** under the optimized conditions (Scheme 3). 2-Methylindole derivatives **1a–f** bearing electron-donating or -withdrawing substituents at C5 gave the corresponding spiroindolenines **2a–f** in high yields with good to moderate enantioselectivities (59–81% ee). The optical purity of **2a–f** could be improved to 88–99%

Scheme 3. Enantioselective Oxidative Aza-spirocyclization to Give Spiropyrrolidines **2** Under the Optimized Conditions^a




^aReaction conditions: **4b** (10 mol %), CHP (2 equiv), Na₂SO₄, MTBE/toluene, 25 °C, 10–24 h. ^bAfter a single recrystallization. ^cReaction conditions: Bu₄NI (10 mol %), CHP (2 equiv), toluene, 25 °C, 10 h.

ee after a single recrystallization. Good to excellent enantioselectivities (75–98% ee) were achieved for the reaction of 2-alkyl (larger than methyl)- or 2-phenyl-substituted indoles bearing electron-withdrawing substituents (**1g–o**, **1q–t**) or no substituent (**1p**) on the indole nucleus. Interestingly, while the reaction of *rac*-**1u** using Bu₄NI gave high diastereoselectivity (12:1), the use of chiral catalyst **4b** afforded a 2:1 mixture of **2u** and **2u'** with higher enantioselectivity for the minor diastereomer **2u'**.

Control experiments revealed that the ammonium hypoiodite species might be the catalytically active species for the oxidative dearomatative spirocyclization of **1** and that a free radical pathway might be unlikely (Table S7).

To gain further insight into the reaction mechanism, we performed kinetic studies using the oxidative dearomatization of **1v**, a *p*-(trifluoromethyl)benzenesulfonyl-protected analogue of **1a**, as a model reaction (Figures S1–S4).²⁰ The reaction rate was found to have a zeroth-order dependence on the concentration of substrate **1v** and a first-order dependence on the concentrations of both CHP and **4b**. However, a difference in the initial reaction rates was observed depending on the substituent of **1**, suggesting that oxidation of a catalyst–substrate complex might be the rate-determining step.²¹ To further evaluate the mechanism, we performed a Hammett analysis with a series of *N*-4-(trifluoromethyl)benzenesulfonyl homotryptamines **1v–y** and *N*-sulfonyl 5-bromohomotryptamines **1z–ab** to probe the electronic effects of the para substituents on the indole and sulfonamide nitrogens, respectively, on the reaction rate (Scheme 4a and Figures S5–S8). As a result, a linear correlation with a negative slope ($\rho = -0.98$) was observed from the corresponding plot of the σ_{para} constants versus $\log(k_{\text{R}}/k_{\text{H}})$ for indole substituents (**R**⁵). On the other hand, a poor correlation was observed from the

Scheme 4. Mechanistic Studies



corresponding plot of the σ_{meta} constants, suggesting the accumulation of positive charge on the indole nitrogen rather than C3 in the rate-determining transition state.^{22,23} Although a linear correlation was also obtained for sulfonamide substituents (R), the reaction constant was much smaller ($\rho = -0.15$), suggesting that accumulation of positive charge on the sulfonamide nitrogen might be unlikely.

Next, to evaluate the roles of the *N*-H groups of the indole and sulfonamide units for the oxidation reactions, *N*-methylindole 5 and *N*-methylsulfonamide 6 were prepared and examined under the standard conditions (Scheme 4b). While most of the starting material was recovered from the reaction of 5, the dearomatization of 6 proceeded smoothly to give peroxide adduct 7 in good yield. Consistent with the results of the Hammett analysis, these results indicated that umpolung of the indole moiety through the generation of an *N*-I indole intermediate might be crucial for the oxidative dearomatization reaction. On the other hand, the smooth reaction of 1d using *N*-iodosuccinimide as a stoichiometric I^+ reagent under neutral conditions gave C3-iodine adduct 8 (Scheme 4c). C3 iodination was not observed under our

catalytic conditions (Figure S9), and exposure of 8 to our conditions gave a complex mixture of several unidentified products; 2d was not observed, suggesting that spirocyclization via iodination of indole C3 might be unlikely.

On the basis of these experimental results and previous works,^{13a,16} a proposed catalytic reaction mechanism is depicted in Scheme 4d. Ammonium hypiodite could be generated *in situ* as an active species from the oxidation of ammonium iodide with an oxidant. *N*-Iodo intermediate 9 might be produced by a reversible reaction of hypiodite with the indole *N*-H directly or by iodination of the sulfonamide *N*-H followed by intramolecular iodo transfer.²⁴ To enhance the electrophilicity²¹ of indole, *N*-iodine(III) intermediate 10 might be generated by rate-determining oxidation of 9. The accumulation of positive charge on the indole nitrogen in the rate-determining transition state with the generation of a highly electron-deficient iodine(III) species is also in agreement with the results of the Hammett analysis. Finally, reductive elimination of ammonium hypiodite might proceed via intramolecular capture of highly electrophilic intermediate 10 by the chiral ammonium sulfonamide as the enantio-determining step to give aza-spiroindolenine 2.


Given the high nucleofugality of hypervalent iodines,²⁵ we considered that zwitterionic intermediate 11 might be generated by the competitive dissociation of ammonium hypiodite prior to spirocyclization,²⁶ which would render asymmetric induction difficult (Scheme 4d). A dissociative racemic pathway might preferentially proceed for the oxidation of electron-rich indoles (e. g., 1e and 1f) as a result of stabilization of cationic intermediate 11 by electron-donating substituents to give moderate enantioselectivity (Scheme 3). In addition, plotting $\log(\text{e.r.})$ ²⁷ of products 2a–f against the corresponding σ_{para} constants gave a linear correlation with a positive slope (Scheme 4d, inset), which might also support the existence of a dissociative pathway during umpolung of indole.

We envisioned that the introduction of an electron-deficient substituent at C2 might suppress the dissociative pathway by destabilization of cationic intermediate 11. In addition, an electron-deficient group at C2 would further reduce the LUMO energy at C3, which might enhance the rate of the spirocyclization step and further improve the chemoselectivity. Moreover, if an electron-deficient auxiliary could be used as a leaving group, the synthetic utility of the products would be enhanced. With these assumptions in mind, we focused on pyrazole as an electron-deficient auxiliary because it can be easily introduced²⁸ and removed via nucleophilic acyl substitution.²⁹ To our delight, spiroindolenines 2ac–ae were obtained after smooth reaction of the corresponding 2-pyrazol-1-ylhomotryptamine derivatives 1ac–ae (Scheme 5a). Most importantly, in sharp contrast to those for 2-methylindole analogues 2a, 2d, and 2f, good enantioselectivities (81–86% ee) were achieved regardless of the electron-donating or -withdrawing substituents at C5. A gram-scale oxidation of 1ac was also achieved with the use of 5 mol % 4b to give 2ac in enantiomerically pure form after recrystallization. In addition, an additive robustness screen analysis³⁰ revealed that a wide range of functional groups were tolerated under our mild conditions, including carbonyls, amines, alkyne, and heteroarenes (Table S5).


In sharp contrast to previous methods,³¹ with the introduction of a pyrazole auxiliary at C2, site-selective spirocyclization of 2-pyrazol-1-yltryptamine 12 at C3 pro-

Scheme 5. Harnessing an Electrophilic Indole by Introducing a Pyrazolyl Auxiliary

a) Enantioselective oxidative spirocyclization of 2-pyrazolylhomotryptamine derivatives



b) Enantioselective oxidative dearomatization of 2-pyrazolyltryptamine derivative




^a4b (5 mol %). ^bAfter a single recrystallization.

ceeded to give the difficult-to-access four-membered spiroazetidine **13** with good enantioselectivity (Scheme 5b).³²


Finally, we demonstrated the synthetic utility of enantioenriched spiroindolenines **2** (Scheme 6). First, the 4-nosyl group

Scheme 6. Transformations


a) Deprotection of nosyl group




b) To α -methylene indolines and subsequent condensation



c) Hydrolysis to oxindole and formal C–C coupling



d) Alcoholysis and azidation



of **2a** could be easily removed under standard deprotection conditions to give free amine **14** (Scheme 6a).¹⁹ Deprotonation of **2a** triggered by N-tosylation gave 2-methyleneindoline **15** in good yield (Scheme 6b, left). Similarly, N-methylation of **2a** followed by condensation with salicylaldehyde afforded spiroindolenine **16**, a structure that is commonly found in photochromic compounds³³ (Scheme 6b, right). On the other hand, the pyrazole auxiliary of **2ac** could be easily removed by Lewis acid-catalyzed hydrolysis to give oxindole **17** quantitatively (Scheme 6c).³⁴ Formal nucleophilic substitution of pyrazole **2ac** with a carbon nucleophile was accomplished by triflation of **17** followed by a Sonogashira coupling reaction to give 2-alkynylspiroindolenine **18**. In addition, ethanolysis^{13a} of pyrazole **2ac** proceeded smoothly to give imino ester **19** in good yield (Scheme 6d, left). Moreover, we found that scandium-catalyzed nucleophilic substitution of pyrazole **2ac** by trimethylsilyl azide followed by intramolecular click cyclization gave tetrazoloindole **20** as a

unique structure, which was confirmed by X-ray analysis (Scheme 6d, right). No loss of enantioselectivity was observed in any of these transformations.

In summary, we have developed an oxidative umpolung strategy for the chiral ammonium hypiodite-catalyzed enantioselective dearomative aza-spirocyclization of homotryptamine derivatives to give the corresponding aza-spiroindolenines with good to excellent enantioselectivity. Mechanistic studies revealed the unusual umpolung reactivity at C3 of indoles by N1 iodination. Moreover, by the introduction of pyrazole as an electron-deficient auxiliary at C2, site-selective spirocyclization of a tryptamine derivative was also achieved to give the difficult-to-access spiroazetidine in an enantioselective manner. Furthermore, to demonstrate the synthetic utility of our dearomative spirocyclization, 2-alkyl- and 2-pyrazole-substituted spiroindolenines were readily converted to various useful and unique structures. These results demonstrate the high potential of hypiodite catalysis for oxidative umpolung of indoles toward the synthesis of polycyclic indole-derived alkaloids.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.2c01852>.

Additional information, synthesis procedures, and spectral data (PDF)

Accession Codes

CCDC 2143581, 2143588–2143591, and 2159841 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, U.K.; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Authors

Muhammet Uyanik – Graduate School of Engineering, Nagoya University, Nagoya 464-8603, Japan; orcid.org/0000-0002-9751-1952; Email: muha@chembio.nagoya-u.ac.jp

Kazuaki Ishihara – Graduate School of Engineering, Nagoya University, Nagoya 464-8603, Japan; orcid.org/0000-0003-4191-3845; Email: ishihara@cc.nagoya-u.ac.jp

Authors

Hiroki Tanaka – Graduate School of Engineering, Nagoya University, Nagoya 464-8603, Japan

Naoya Ukegawa – Graduate School of Engineering, Nagoya University, Nagoya 464-8603, Japan

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacs.2c01852>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Financial support for this project was partially provided by JSPS KAKENHI (20K20559 to K.I., 21H01932 to M.U., and 18H01973 to M.U.), the Society of Iodine Science (to M.U.), and the Nagoya University Graduate Program of Trans-

formative Chem-Bio Research (GTR). This work is partially supported by Nagoya University Research Fund.

REFERENCES

- (1) (a) Seigler, D. S. *Indole Alkaloids*. In *Plant Secondary Metabolism*; Springer: Boston, MA, 1998; pp 628–654. (b) Ban, Y.; Murakami, Y.; Iwasawa, Y.; Tsuchiya, M.; Takano, N. Indole alkaloids in medicine. *Med. Res. Rev.* **1988**, *8*, 231–308. (c) de Sá Alves, F. R.; Barreiro, E. J.; Manssour Fraga, C. A. From nature to drug discovery: The indole scaffold as a 'privileged structure'. *Mini-Rev. Med. Chem.* **2009**, *9*, 782–793.
- (2) Ishikura, M.; Abe, T.; Choshi, T.; Hibino, S. Simple indole alkaloids and those with a nonrearranged monoterpenoid unit. *Nat. Prod. Rep.* **2015**, *32*, 1389–1471.
- (3) (a) Roche, S. P.; Tendoung, J.-J. Y.; Tréguier, B. Advances in dearomatization strategies of indoles. *Tetrahedron* **2015**, *71*, 3549–3591. (b) Saya, J. M.; Ruijter, E.; Orru, R. V. A. Total synthesis of aspidosperma and strychnos alkaloids through indole dearomatization. *Chem. - Eur. J.* **2019**, *25*, 8916–8935. (c) Zheng, C.; You, S.-L. Catalytic asymmetric dearomatization (CADA) reaction-enabled total synthesis of indole-based natural products. *Nat. Prod. Rep.* **2019**, *36*, 1589–1605. (d) Sheng, F.-T.; Wang, J.-Y.; Tan, W.; Zhang, Y.-C.; Shi, F. Progresses in organocatalytic asymmetric dearomatization reactions of indole derivatives. *Org. Chem. Front.* **2020**, *7*, 3967–3998. (e) Xia, Z.-L.; Xu-Xu, Q.-F.; Zheng, C.; You, S.-L. Chiral phosphoric acid-catalyzed asymmetric dearomatization reactions. *Chem. Soc. Rev.* **2020**, *49*, 286–300. (f) Zheng, C.; You, S.-L. Advances in Catalytic Asymmetric Dearomatization. *ACS Cent. Sci.* **2021**, *7*, 432–444.
- (4) (a) Bartoli, G.; Bencivenni, G.; Dalpozzo, R. Organocatalytic strategies for the asymmetric functionalization of indoles. *Chem. Soc. Rev.* **2010**, *39*, 4449–4465. (b) Lakhdar, S.; Westermaier, M.; Terrier, F.; Goumont, R.; Boubaker, T.; Ofial, A. R.; Mayr, H. Nucleophilic reactivities of indoles. *J. Org. Chem.* **2006**, *71*, 9088–9095.
- (5) (a) Bandini, M. Electrophilicity: the "dark-side" of indole chemistry. *Org. Biomol. Chem.* **2013**, *11*, 5206–5212. (b) Abou-Hamdan, H.; Kouklovsky, C.; Vincent, G. Dearomatization reactions of indoles to access 3D indoline structures. *Synlett* **2020**, *31*, 1775–1788.
- (6) Rkein, B.; Bigot, A.; Birbaum, L.; Manneveau, M.; De Paolis, M.; Legros, J.; Chataigner, I. Reactivity of 3-nitroindoles with electron-rich species. *Chem. Commun.* **2021**, *57*, 27–44.
- (7) (a) Nagayoshi, T.; Saeki, S.; Hamana, M. Studies on tertiary amine oxides. LXXII. Some nucleophilic reactions of 1-hydroxy-2-phenylindole. *Chem. Pharm. Bull.* **1981**, *29*, 1920–1926. (b) Li, C.; Zhang, H.-H.; Fan, T.; Shen, Y.; Wu, Q.; Shi, F. Brønsted Acid-Catalyzed Regioselective Reactions of 2-Indolylmethanols with Cyclic Enaminone and Anhydride Leading to C3-Functionalized Indole Derivatives. *Org. Biomol. Chem.* **2016**, *14*, 6932–6936. (c) Zhang, H.-H.; Wang, C.-S.; Li, C.; Mei, G.-J.; Li, Y.; Shi, F. Design and Enantioselective Construction of Axially Chiral Naphthyl-Indole Skeletons. *Angew. Chem., Int. Ed.* **2017**, *56*, 116–121. (d) Zhang, Y.-C.; Jiang, F.; Shi, F. Organocatalytic Asymmetric Synthesis of Indole-Based Chiral Heterocycles: Strategies, Reactions, and Outreach. *Acc. Chem. Res.* **2020**, *53*, 425–446. (e) Petrini, M. New Perspectives in the Indole Ring Functionalization using 2-Indolylmethanols. *Adv. Synth. Catal.* **2020**, *362*, 1214–1232.
- (8) Seebach, D. Methods of Reactivity Umpolung. *Angew. Chem., Int. Ed.* **1979**, *18*, 239–258.
- (9) (a) Awang, D. V. C.; Vincent, A. Alkoxylation of 2,3-dialkylindoles utilizing iodosobenzene diacetate. *Can. J. Chem.* **1980**, *58*, 1589–1591. (b) Dohi, T.; Ito, M.; Yamaoka, N.; Morimoto, K.; Fujioka, H.; Kita, Y. Hypervalent iodine(III): selective and efficient single-electron-transfer (SET) oxidizing agent. *Tetrahedron* **2009**, *65*, 10797–10815. (c) Caramenti, P.; Waser, J. Bench-stable electrophilic indole and pyrrole reagents: serendipitous discovery and use in C–H functionalization. *Helv. Chim. Acta* **2017**, *100*, e1700221.
- (10) (a) Wu, J.; Dou, Y.; Guillot, R.; Kouklovsky, C.; Vincent, G. Electrochemical dearomative 2,3-difunctionalization of indoles. *J. Am. Chem. Soc.* **2019**, *141*, 2832–2837. (b) Liu, K.; Song, W.; Deng, Y.; Yang, H.; Song, C.; Abdelilah, T.; Wang, S.; Cong, H.; Tang, S.; Lei, A. Electrooxidation enables highly regioselective dearomative annulation of indole and benzofuran derivatives. *Nat. Commun.* **2020**, *11*, 3.
- (11) (a) Gentry, E. C.; Rono, L. J.; Hale, M. E.; Matsuura, R.; Knowles, R. R. Enantioselective synthesis of pyrroloindolines via noncovalent stabilization of indole radical cations and applications to the synthesis of alkaloid natural products. *J. Am. Chem. Soc.* **2018**, *140*, 3394–3402. (b) Cheng, Y.-Z.; Zhao, Q.-R.; Zhang, X.; You, S.-L. Asymmetric dearomatization of indole derivatives with N-hydroxycarbamates enabled by photoredox catalysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 18069–18074. (c) Yuan, W.; Huang, J.; Xu, X.; Wang, L.; Tang, X.-Y. B(C₆F₅)₃-Catalyzed electron donor-acceptor complex-mediated aerobic sulfenylation of indoles under visible-light conditions. *Org. Lett.* **2021**, *23*, 7139–7143. (d) Cheng, Y.-Z.; Feng, Z.; Zhang, X.; You, S.-L. Visible-light Induced Dearomatization Reactions. *Chem. Soc. Rev.* **2022**, *51*, 2145.
- (12) (a) Uyanik, M.; Okamoto, H.; Yasui, T.; Ishihara, K. Quaternary ammonium (hypo)iodite catalysis for enantioselective oxidative cycloetherification. *Science* **2010**, *328*, 1376–1379. (b) Uyanik, M.; Ishihara, K. Catalysis with *in situ* generated (hypo)iodite ions for oxidative coupling reactions. *ChemCatChem* **2012**, *4*, 177–185. (c) Finkbeiner, P.; Nachtsheim, B. J. Iodine in modern oxidation catalysis. *Synthesis* **2013**, *45*, 979–999. (d) Chen, R.; Chen, J.; Zhang, J.; Wan, X. Combination of tetrabutylammonium iodide (TBAI) with *tert*-butyl hydroperoxide (TBHP): an efficient transition-metal-free system to construct various chemical bonds. *Chem. Rec.* **2018**, *18*, 1292–1305.
- (13) (a) Uyanik, M.; Tanaka, H.; Ishihara, K. Hypoiodite-catalyzed chemoselective tandem oxidation of homotryptamines to peroxy- and epoxytetrahydropyridindolenines. *Org. Lett.* **2020**, *22*, 8049–8054. (b) Uyanik, M.; Tanaka, H.; Ishihara, K. I⁺/TBHP catalysis for tandem oxidative cyclization to indolo[2,3-*b*]quinolines. *Asian J. Org. Chem.* **2021**, *10*, 164–169.
- (14) Zheng, C.; You, S.-L. Exploring the chemistry of spiroindolenines by mechanistically-driven reaction development: Asymmetric Pictet–Spengler-type reactions and beyond. *Acc. Chem. Res.* **2020**, *53*, 974–987.
- (15) (a) Tanaka, K.; Mori, Y.; Narasaka, K. Synthesis of spiro[indoline-3,2'-pyrrolidine] derivatives from β-3-indolyl ketone oximes. *Chem. Lett.* **2004**, *33*, 26–27. (b) Ma, X.; Farndon, J. J.; Young, T. A.; Fey, N.; Bower, J. F. A simple and broadly applicable C–N bond forming dearomatization protocol enabled by bifunctional amino-reagents. *Angew. Chem., Int. Ed.* **2017**, *56*, 14531–14535. (c) Wang, P.-F.; Chen, C.; Chen, H.; Han, L.-S.; Liu, L.; Sun, H.; Wen, X.; Xu, Q.-L. Concise synthesis of spiro[indoline-3,2'-pyrrolidine] and 1-azacarbazole derivatives via copper-catalyzed cyclization of indoles. *Adv. Synth. Catal.* **2017**, *359*, 2339–2344. (d) Zhou, Y.; Li, D.; Tang, S.; Sun, H.; Huang, J.; Zhu, Q. PhI(OAc)₂-mediated dearomative C–N coupling: facile construction of the spiro[indoline-3,2'-pyrrolidine] skeleton. *Org. Biomol. Chem.* **2018**, *16*, 2039–2042.
- (16) Uyanik, M.; Hayashi, H.; Ishihara, K. High-turnover hypoiodite catalysis for asymmetric synthesis of tocopherols. *Science* **2014**, *345*, 291–294.
- (17) Almost no reaction was observed with the use of hydrogen peroxide as an oxidant, suggesting nonproductive decomposition of the oxidant. See: Uyanik, M.; Sahara, N.; Tsukahara, M.; Hattori, Y.; Ishihara, K. Chemo- and enantioselective oxidative α-azidation of carbonyl compounds. *Angew. Chem., Int. Ed.* **2020**, *59*, 17110–17117.
- (18) Ooi, T.; Maruoka, K. Recent advances in asymmetric phase-transfer catalysis. *Angew. Chem., Int. Ed.* **2007**, *46*, 4222–4266.
- (19) Considering further synthetic study, we used a *p*-nitrobenzenesulfonyl (^tNs) group as a useful protecting group for easy deprotection (Table S2). See: Kan, T.; Fukuyama, T. Ns strategies: a highly versatile synthetic method for amines. *Chem. Commun.* **2004**, 353–359.

(20) Considering the better solubility of substrates **1** under the reaction conditions, we selected a (trifluoromethyl)benzenesulfonyl group as a protecting group for kinetic and Hammett studies.

(21) Bosnidou, A. E.; Muñiz, K. Alkyl iodides in high oxidation state: enhanced synthetic possibilities and accelerated catalyst turn-over. *Chem. - Eur. J.* **2019**, *25*, 13654–13664.

(22) Richter, J. M.; Whitefield, B. W.; Maimone, T. J.; Lin, D. W.; Castroviejo, M. P.; Baran, P. S. Scope and mechanism of direct indole and pyrrole couplings adjacent to carbonyl compounds: total synthesis of acremoauxin A and oxazinin 3. *J. Am. Chem. Soc.* **2007**, *129*, 12857–12869.

(23) Nandi, R. K.; Perez-Luna, A.; Gori, D.; Beaud, R.; Guillot, R.; Kouklovsky, C.; Gandon, V.; Vincent, G. Triflic Acid as an Efficient Brønsted Acid Promoter for the Umpolung of *N*-Ac Indoles in Hydroarylation Reactions. *Adv. Synth. Catal.* **2018**, *360*, 161–172.

(24) Although we could not isolate and fully characterize *N*-iodo intermediate **9**, possibly because of inherent instability of the compound and rapid equilibrium with **1**, NMR and mass analysis of the reaction mixture suggested iodination of the indole to give a transient intermediate (Figures S9 and S10).

(25) Yoshimura, A.; Zhdankin, V. V. Advances in synthetic applications of hypervalent iodine compounds. *Chem. Rev.* **2016**, *116*, 3328–3435.

(26) Uyanik, M.; Nishioka, K.; Kondo, R.; Ishihara, K. Chemo-selective oxidative generation of *ortho*-quinone methides and tandem transformations. *Nat. Chem.* **2020**, *12*, 353–362.

(27) Wendlandt, A. E.; Vangal, P.; Jacobsen, E. N. Quaternary stereocentres via an enantioconvergent catalytic S_N1 reaction. *Nature* **2018**, *556*, 447–451.

(28) Wu, W.-B.; Huang, J.-M. Highly regioselective C–N bond formation through C–H azolation of indoles promoted by iodine in aqueous media. *Org. Lett.* **2012**, *14*, 5832–5835.

(29) Tokumasu, K.; Yazaki, R.; Ohshima, T. Direct catalytic chemoselective α -amination of acylpyrazoles: A concise route to unnatural α -amino acid derivatives. *J. Am. Chem. Soc.* **2016**, *138*, 2664–2669.

(30) Collins, K.; Glorius, F. A Robustness Screen for the Rapid Assessment of Chemical Reactions. *Nat. Chem.* **2013**, *5*, 597–601.

(31) Hexahydropyrroloindole structures were often obtained via cyclization of tryptamine at C2. See: Ruiz-Sanchis, P.; Savina, S. A.; Albericio, F.; Alvarez, M. Structure, bioactivity and synthesis of natural products with hexahydropyrrolo[2,3-*b*]indole. *Chem. - Eur. J.* **2011**, *17*, 1388–1408.

(32) Although there remains room for improvement in the enantioselectivity, preliminary experiments demonstrated the prospects of the use of the 2-pyrazole auxiliary to expand the reaction scope to spiroaziridination and spiro-lactonization. See Scheme S1.

(33) Kortekaas, L.; Browne, W. R. The evolution of spiroopyran: fundamentals and progress of an extraordinarily versatile photochrome. *Chem. Soc. Rev.* **2019**, *48*, 3406–3424.

(34) Quite recently, Zhong and colleagues reported the enantioselective synthesis of spiroindolyloxindoles as related compounds by chiral ammonium hypoiodite-catalyzed oxidative α -aminocyclization of carbonyl compounds. See: Wang, D.; Zhang, W.; Lu, X.; Zhou, H.; Zhong, F. Cinchona Alkaloid Derived Iodide Catalyzed Enantioselective Oxidative α -Amination of Carbonyl Compounds toward the Construction of Spiroindolyloxindole. *Org. Lett.* **2022**, *24*, 842–847.