


Highly Efficient Enantioselective Synthesis of Chiral Sulfones by Rh-Catalyzed Asymmetric Hydrogenation

Qiaozhi Yan,[†] Guiying Xiao,[†] Ying Wang, Guofu Zi,[‡] Zhanbin Zhang, and Guohua Hou*[‡]

Key Laboratory of Radiopharmaceuticals, College of Chemistry, Beijing Normal University, Beijing 100875, China

S Supporting Information

ABSTRACT: A highly efficient and enantioselective Rh-(*R,R*)-*f*-spiroPhos complex catalyzed hydrogenation of a series of unsaturated sulfones has been developed. With Rh-(*R,R*)-*f*-spiroPhos catalyst under mild conditions, not only the asymmetric hydrogenation of both the 3,3-diaryl and exocyclic α,β -unsaturated sulfones was first realized with up to 99.9% ee but also 3-alkyl-3-aryl and benzo[*b*]thiophene-1,1-dioxides were successfully hydrogenated to the corresponding chiral sulfones with excellent enantioselectivities (up to 99.4% ee) regardless of the steric hindrance, electronic property, and geometry of the substrates. Moreover, this reaction offers a route to (*S*)-(+)-*ar*-turmerone as a spice flavor, which is an important synthetic intermediate of pharmaceuticals.



1. INTRODUCTION

Chiral sulfones represent an important functional group prevalently found in many biologically active compounds and pharmaceutical intermediates to produce *ar*-turmerone,¹ (*R*)-curcuphenol,^{1b} (*S*)-curcudiol,^{1b} Remikiren,² *r*-secretase inhibitor,³ and inflammatory cytokines inhibitor (Figure 1).⁴ Sulfones serve as useful intermediates and can be converted into other chiral functional groups by alkylation, halogenation, oxidation, desulfonylation, and Julia olefinations.⁵

The synthesis of chiral sulfones has attracted increased attention due to their utility as intermediates. Previous reported methods for the synthesis of chiral sulfones have




Figure 1. Key structural elements in chiral pharmaceuticals.

included asymmetric arylations/alkenylations of α -bromosulfones,⁶ rhodium- and organo-catalyzed asymmetric conjugate addition to unsaturated sulfones,⁷ asymmetric radical additions,⁸ and asymmetric conjugate reduction of prochiral unsaturated sulfones.^{9–11,13–16} Among the reported methods, metal-catalyzed asymmetric conjugate reduction of prochiral unsaturated sulfones is of interest as one of the most straightforward approaches for chiral sulfone synthesis. However, only a few efficient catalyst systems for the preparation of chiral sulfones have been reported so far.^{10,11}

Catalytic asymmetric hydrogenation represents one of the most efficient, straightforward, and well-established approaches to chiral compounds.¹² To date, only a few examples of asymmetric hydrogenation of unsaturated sulfones have been reported.^{13–17} Andersson and co-workers reported the efficient asymmetric hydrogenation of vinylic, allylic, homoallylic, and cyclic sulfones catalyzed by a chiral Ir/N,P complex achieving good to high enantioselectivities.¹³ Recently, Tosatti and Pfaltz also developed two chiral iridium N,P-ligand complexes that exhibited good activity and enantioselectivity in the asymmetric hydrogenation of benzothiophene 1,1-dioxides. However, only moderate to good enantioselectivities were achieved in the hydrogenation of the corresponding alkyl substituted substrates.¹⁴ Zhang and co-workers developed Rh-catalyzed asymmetric hydrogenation of α -substituted vinyl sulfones and β -acetylamino acrylosulfones using DTBM-SegPhos and Tangphos, respectively, achieving high enantioselectivities for the (*Z*)- β -acetylamino acrylosulfones.^{15,16} However, (*E*)-substrates only provided low to moderate enantioselectivities.

Received: November 26, 2018

Published: January 7, 2019

Thus, the substrate scope of the asymmetric hydrogenation of sulfones remains limited.

Representative challenging sulfone-containing moieties for asymmetric hydrogenation are shown in Figure 2. The





Figure 2. Some challenging substrates of unsaturated sulfones.

following sterically hindered substrates including 3,3-diaryl unsaturated sulfones and the cyclic unsaturated sulfones have so far remained unexplored as substrates for asymmetric hydrogenation. It is not surprising because the bigger aromatic system and the bulkier steric hindrance in the substrates results in their inertness, which requires much higher catalytic activity. So far only a few successful examples of asymmetric hydrogenation of the substrates bearing diaryl substituents at the stereogenic center were reported.¹⁸ In addition, most of the present catalysts are only efficient for either of two isomers of unsaturated sulfones. Therefore, the development of efficient catalysts for asymmetric hydrogenation of unsaturated sulfones to afford chiral sulfones is still highly desirable.

Recently, we prepared and demonstrated the chiral diphosphine ligand, *f*-spiroPhos, containing the 1,1'-spirobiindane skeleton, as a highly efficient and enantioselective ligand in the hydrogenation of various substrates including unsaturated nitriles, carboxylic acids, and imines.¹⁹ Encouraged by the previous work, we decided to attempt to tackle the asymmetric hydrogenation of these challenging unsaturated sulfones. We herein report the first example of highly enantioselective Rh-catalyzed hydrogenation of 3,3-diaryl and exocyclic unsaturated sulfones with up to 99.9% ee. The reaction also showed high substrate scope; *E*- and *Z*-isomers of a wide range of substrates including 3,3-diaryl, 3-alkyl-3-aryl, exocyclic α,β -unsaturated sulfones, and benzo[*b*]thiophene 1,1-dioxide can also be hydrogenated with excellent enantioselectivities under mild reaction conditions (Scheme 1).

Scheme 1



2. RESULTS AND DISCUSSION

2.1. Asymmetric Hydrogenation of Diaryl α,β -Unsaturated Sulfones. Our initial investigation began with the hydrogenation of (*E*)-1-chloro-3-(1-phenyl-2-(phenylsulfonyl)vinyl)benzene **1a** as the model substrate under 80 atm of H₂ in CH₂Cl₂ at room temperature using 1.0 mol % of the complex of [Rh(COD)₂]BF₄ and (*R,R*)-*f*-spiroPhos. Although only a moderate conversion was provided,

Table 1. Asymmetric Hydrogenation of (*E*)-1-Chloro-3-(1-phenyl-2-(phenylsulfonyl)vinyl)benzene **1a**^a

entry	ligand	solvent	conv ^b (%)	ee ^c (%)
1	(<i>R,R</i>)- <i>f</i> -spiroPhos	CH ₂ Cl ₂	60	96
2	(<i>R,R</i>)- <i>f</i> -spiroPhos	MeOH	5	ND
3	(<i>R,R</i>)- <i>f</i> -spiroPhos	THF	3	ND
4	(<i>R,R</i>)- <i>f</i> -spiroPhos	DCE	6	ND
5	(<i>R,R</i>)- <i>f</i> -spiroPhos	dioxane	3	ND
6	(<i>R,R</i>)- <i>f</i> -spiroPhos	toluene	99	98
7	(<i>S</i>)-BINAP	toluene	0.5	ND
8	(<i>R</i>)-JosiPhos-1	toluene	2	ND
9	(<i>S,S</i>)-Me-Duphos	toluene	1	ND
10	(<i>R,R</i>)-Ph-BPE	toluene	1	ND
11	(<i>S,S</i>)- <i>f</i> -Binaphane	toluene	29	68

^aReaction conditions: Rh(COD)₂BF₄/ligand/substrate ratio = 1.0:1.1:100, 80 atm H₂, 24 h, rt. ^bDetermined by GC analysis. ^cDetermined by chiral supercritical fluid chromatography (SFC) using a chiral stationary phase Lux 5u Cellulose-3 (250 × 4.60 mm), MeOH/CO₂ = 30:70, 3.0 mL/min, 210 nm.

a high enantioselectivity was achieved, 96% ee (Table 1, entry 1). To increase substrate conversion, we subsequently investigated the solvent effect. The role of solvent was found to play an important role in the asymmetric hydrogenation of chiral sulfones. Poor conversion was observed in most of the solvents such as MeOH, tetrahydrofuran (THF), dichloroethane (DCE), and 1,4-dioxane (Table 1, entries 2–5). To our delight, by performing reactions in toluene as the solvent, complete conversion and high enantioselectivity (98% ee) were achieved (Table 1, entry 6). Following solvent optimization, a screening of commercially available chiral phosphorus ligands (Figure 3) revealed that (*S*)-BINAP, electron-rich (*R*)-JosiPhos, (*S,S*)-Me-Duphos, and (*R,R*)-Ph-BPE were ineffective in this transformation and only trace products were afforded (Table 1, entries 7–10). (*S,S*)-*f*-Binaphane gave higher conversion (29%) and ee value (68%), thus demonstrating that the rigidity and bulkier steric hindrance of the spiro-backbone in *f*-spiroPhos had a significant positive effect on the conversion and enantioselectivity in this hydrogenation.

Encouraged by the promising results obtained by the asymmetric hydrogenation of (*E*)-1-chloro-3-(1-phenyl-2-(phenylsulfonyl)vinyl)benzene **1a**, we then prepared a variety of 3,3-diaryl unsaturated sulfones **1** and applied them to the hydrogenation in toluene under the catalyst loading of 1.0 mol % [Rh(COD)₂]BF₄/*(R,R)*-*f*-spiroPhos and 80 atm H₂ (Table 2). All of the substrates examined (**1a–1o**) could be successfully hydrogenated to produce the corresponding chiral sulfones in high yields with excellent enantioselectivities, 94–99.9% ee. Neither the electron property nor the position of the substituents at the two aryl rings of the substrates had an apparent effect on the enantioselectivity. In general, the substrates with electron-donating substituents including Me and MeO group exhibited relatively lower reactivity compared with those bearing electron-withdrawing substituents. For instance, the fluoro- or chloro-substituted substrates **1a–1g** were hydrogenated in full conversions with 96–99.9% ee,





Figure 3. Structures of the phosphine ligands for hydrogenation of (*E*)-1-chloro-3-(1-phenyl-2-(phenylsulfonyl)vinyl)benzene **1a**.

Table 2. Rh-Catalyzed Asymmetric Hydrogenation of 3,3-Diaryl α,β -Unsaturated Sulfones **1**^a



^aUnless otherwise mentioned, all reactions were carried out with a [Rh(COD)₂]₂BF₄/(*R,R*)-f-spiroPhos/substrate ratio of 1.0:1.1:100, toluene, 80 atm H₂, rt, 24 h. The conversion was determined by ¹H NMR spectroscopy or GC analysis; the enantioselectivity was determined by SFC analysis using a chiral stationary phase.

while the substrates bearing electron-donating substituents Me and MeO **1h–1l** provided 94–98% conversions. Notably, even if the substrate contained the same substituent present at different positions of the two-phenyl ring (i.e., meta and para), excellent enantioselectivity was achieved. This is shown in the following example: substrates **1i** and **1n** substituted by a methyl group at the meta-position of one phenyl ring or both meta- and para-position of the two phenyl rings, respectively, provided the highest enantioselectivity, 99.9% ee. Remarkably, this catalyst could also exhibit comparable activity and enantioselectivity in the hydrogenation of the corresponding (*Z*)-isomers to afford the products with an opposite absolute configuration compared with that obtained by (*E*)-isomers. For example, the substrates (*Z*)-**1b** and (*Z*)-**1d** were hydrogenated, providing the desired products **2b** and **2d** with 99.9% and 98% ee, respectively. It was found that the (*Z*)-substrates provided the hydrogenation products with an opposite absolute configuration compared with that afforded by (*E*)-isomers by comparison with the sign of the specific rotation and order of elution for major and minor enantiomers from the chiral column. The absolute configuration of hydrogenation product **2l** was determined and assigned to be the (*R*) configuration by X-ray crystallographic analysis of the corresponding single-crystal structure (Figure 4). It was demonstrated that this method could produce chiral sulfones with either desired configuration, (*S*)- or (*R*)-enantiomer, in excellent enantioselectivities by the asymmetric hydrogenation of (*E*)- or (*Z*)- α,β -unsaturated sulfones.

2.2. Asymmetric Hydrogenation of Cyclic α,β -Unsaturated Sulfones. Previously, Desrosiers and Charette reported the copper-DuPhos(O)-catalyzed enantioselective hydrosilylation of vinyl phenyl sulfones including two cyclic sulfones and achieved excellent enantioselectivities.¹¹ Surprisingly, the asymmetric hydrogenation of cyclic unsaturated sulfones (substrate **3**) has not been reported. To develop the asymmetric hydrogenation of this class of substrates and extend the substrate scope of the catalyst, we decided to evaluate the cyclic α,β -unsaturated sulfone **3a** under the previously optimized reaction conditions in the hydrogenation of 3,3-diaryl α,β -unsaturated sulfones **1**; a conversion of 76% and 90% ee were obtained. After further solvent optimization, dichloromethane was found to result in full conversion and better enantioselectivity (93% ee). Lowering the hydrogen






Figure 4. X-ray crystal structure of product (R)-2I.

Table 3. Rh-Catalyzed Asymmetric Hydrogenation of Cyclic α,β -Unsaturated Sulfones **3**^a

^aUnless otherwise mentioned, all reactions were carried out with a $[\text{Rh}(\text{COD})_2]\text{BF}_4/(\text{R},\text{R})\text{-f-spiroPhos}$ /substrate ratio of 1.0:1.1:100, CH_2Cl_2 , 10 atm H_2 , rt, 1 h. The conversion was determined by ^1H NMR spectroscopy; the enantioselectivity was determined by high-performance liquid chromatography (HPLC) analysis using a chiral stationary. ^b80 atm H_2 , 24 h. ^c $[\text{Rh}(\text{COD})_2]\text{BF}_4$ (2.0 mol %), 80 atm H_2 , 24 h. ^d $[\text{Rh}(\text{COD})_2]\text{BF}_4$ (3 mol %), 80 atm H_2 , 24 h.


pressure from 80 to 10 atm still resulted in complete substrate conversion while preserving high enantioselectivity. The effect of different electronic substitutes on the ring was investigated, and all resulted in full conversions and excellent enantioselectivities (>92% ee). It was revealed that ring size in the substrates (i.e., cyclopentyl, cyclohexyl, or cycloheptyl) had no apparent influence on the enantioselectivity (substrates **3a–3h**), and excellent enantioselectivities were observed (Table 3). However, the reactivity was affected by the ring size of the substrates. The substrates with five-membered ring **3a–3d** could be hydrogenated to obtain the corresponding products in satisfactory conversions under 10 atm of H_2 at room temperature for 1 h, whereas the substrate with a six- or seven-membered ring required higher catalyst loadings and increased

Table 4. Rh-Catalyzed Asymmetric Hydrogenation of Benzo[*b*]thiophene 1,1-Dioxides **5**^a

^aUnless otherwise mentioned, all reactions were carried out with a $[\text{Rh}(\text{COD})_2]\text{BF}_4/(\text{R},\text{R})\text{-f-spiroPhos}$ /substrate ratio of 1.0:1.1:100, CH_2Cl_2 , 10 atm H_2 , rt, 1 h. The conversion was determined by ^1H NMR spectroscopy; the enantioselectivity was determined by HPLC analysis using a chiral stationary. ^b80 atm H_2 , 24 h.

hydrogen pressure for complete conversion. The substrate (*E*)-**3f** could be fully hydrogenated under 80 atm of hydrogen pressure in 24 h, while (*E*)-**3g** with a seven-membered ring required 2 mol % catalyst loading. The catalyst system was also suitable for the (*Z*)-isomer of the cyclic unsaturated sulfones, such as the (*Z*)-**3h**, providing the corresponding product **4h** with excellent enantioselectivities (93% ee), with a reversal of the absolute configuration as discussed earlier.

2.3. Asymmetric Hydrogenation of Benzo[*b*]thiophene 1,1-Dioxides. Recently, Pfaltz's group reported Ir–P,N complex catalyzed asymmetric hydrogenation of benzo[*b*]thiophene 1,1-dioxides, and the 2- and 3-aryl substituted substrates were hydrogenated with high conversions and excellent enantioselectivities (up to 99% ee).¹⁴ However, poor conversions or moderate to good enantioselectivities were observed for the 2-alkyl-substituted substrates. The application of the catalyst/ligand investigated reported herein, Rh-(*R,R*)-f-spiroPhos, for the hydrogenation of 2-alkyl-

Table 5. Rh-Catalyzed Asymmetric Hydrogenation of (*E*)-3-Alkyl-3-aryl α,β -Unsaturated Sulfones 7^a

^aUnless otherwise mentioned, all reactions were carried out with a [Rh(COD)₂]BF₄/(*R,R*)-f-spiroPhos/substrate ratio of 1.0:1.1:100, CH₂Cl₂, 10 atm H₂, rt, 1 h. The conversion was determined by ¹H NMR spectroscopy or GC analysis; the enantioselectivity was determined by HPLC analysis using a chiral stationary phase. ^b10 atm H₂, 2 h, 1,4-dioxane. ^cRh(COD)₂PF₆ (2 mol %), 80 atm H₂, 24 h.





Figure 5. Substrates evaluated on a large scale.


Scheme 2



substituted benzo[*b*]thiophene 1,1-dioxides yielded excellent enantioselectivities (up to 99% ee) for a wide range of 2-alkyl-substituted substrates 5 (Table 4). The 2-methyl-, 2-ethyl-, and

2-*n*-propyl-substituted substrates 5a–5c were hydrogenated under 10 atm hydrogen pressure in 1 h to provide the products with high enantioselectivities and conversions. Replacing the alkyl substituent at 2-position with an isopropyl group ((*E*)-5d) still maintained the excellent enantioselectivity of 96% ee but led to a decreased conversion of 81%, and this was attributed to the increased steric hindrance. The 2-phenyl-substituted substrate (*E*)-5f also exhibited lower reactivity, but higher hydrogen pressure and a longer reaction time for satisfactory conversion were required. Inspired by these exciting results obtained by these cyclic substrates, we also investigated the hydrogenation of linear α -methyl- or phenyl-substituted substrates 5g or 5h, but very poor conversions (5% and 20% conversion, respectively) were observed.

2.4. Asymmetric Hydrogenation of 3-Alkyl-3-aryl α,β -Unsaturated Sulfones. Finally, we turned our attention to the asymmetric hydrogenation of 3-alkyl-3-aryl α,β -unsaturated sulfones. Initially, we chose (*E*)-((2-phenylprop-1-en-1-yl)-sulfonyl)benzene 7a as the model substrate under 80 atm of H₂

Scheme 3. Proposed Enantiodetermining Transition Model for Rh-Catalyzed Asymmetric Hydrogenation of α,β -Unsaturated Sulfones

in CH_2Cl_2 . Both full conversion and high enantioselectivity of 94% ee were obtained. For comparison, other electron-rich ligands including (*R*)-JosiPhos-1, (*S,S*)-Me-Duphos, (*R,R*)-Ph-BPE, and (*S,S*)-f-Binaphane were also evaluated, but only poor conversions and enantioselectivities were obtained except for the ligand (*S,S*)-f-Binaphane, which achieved relatively better results, 71% conversion and 77% ee.²⁰ As shown earlier, the hydrogen pressure can be lowered to 10 atm without any deleterious effects on enantioselectivity or reaction time (1 h). When further lowering the hydrogen pressure to 5 atm, high conversion (97%) and high enantioselectivity were achieved. In addition to CH_2Cl_2 , 1,4-dioxane was also proved as a suitable solvent for this hydrogenation, and an extremely high enantioselectivity of 99% ee with full conversion could be achieved.

It was revealed that a variety of (*E*)-3-alkyl-3-aryl α,β -unsaturated sulfones **7** could be successfully hydrogenated to produce the corresponding chiral sulfones with 93–99.4% ee (Table 5) using the catalyst system. The electron property and position of the substituents at the phenyl ring of the substrates had no obvious influence on both the reactivity and enantioselectivity. Generally, the substrates with electron-donating substituents on the phenyl ring gave higher enantioselectivities than those with electron-withdrawing substituents. For example, substrates (*E*)-**7i** and (*E*)-**7j** with a *para*-fluoro or *para*-chloro substituent gave 93–94% ee, while substrate (*E*)-**7k** with a *para*-methyl substituent yielded **8k** with as high as 99% ee. The substrates with a larger 3-alkyl substituent such as an ethyl group ((*E*)-**7l**) could also give comparable enantioselectivity with substrate (*E*)-**7a**. Both the 1-naphthyl substrate (*E*)-**7m** and 2-naphthyl (*E*)-**7n** exhibited higher enantioselectivity, and (*E*)-**7n** provided product **8n** with the highest 99.4% ee value. The hydrogenation of the (*Z*)-isomers such as the substrates (*Z*)-**7o** and (*Z*)-**7p** could also be performed smoothly, providing the desired products **8o** and **8p** with 97% and 96% ee, respectively, which was the highest enantioselectivity in the hydrogenation of the (*Z*)-isomers of 3-alkyl-3-aryl α,β -unsaturated sulfones so far. It was also found that the products obtained by the (*E*)- and (*Z*)-isomers were opposite enantiomers.

2.5. Preparative Asymmetric Hydrogenation on a Larger Scale of Unsaturated Sulfones. Under the optimal reaction conditions, we carried out the preparative hydrogenation of the following substrates in Figure 5 on a large scale (1.0 mmol). The desired products were obtained with maintained high conversions and excellent enantioselectivities.

2.6. Application of the Rh-(*R,R*)-f-spiroPhos Catalyst to the Synthesis of Biologically Active Compounds and Pharmaceuticals. The optical active chiral sulfones are

versatile synthetic intermediates that can be easily converted to various derivatives. To illustrate this point, we performed a short synthesis of the precursor of (*S*)-(+)-*ar*-turmerone and (+)-bisacumol (Scheme 2). Product (*R*)-**8k** obtained by asymmetric hydrogenation could be transformed by acylation reaction with 3,3-dimethylacrylic acid methyl ester and $\text{MgBr}_2 \cdot \text{Et}_2\text{O}$ to intermediate **9**. In the following step, intermediate **9** was converted to (*S*)-(+)-*ar*-turmerone as a spice flavor of turmeric. (*S*)-(+)-*ar*-Turmerone could also be used for the asymmetric synthesis of (*S*)-*ar*-himachalene, which is a pheromone component of the flea beetle and (+)-bisacumol, as a member of the aromatic bisabolane sesquiterpene family isolated from the rhizome of *Curcuma Xanthorrhiza*.^{1b}

On the basis of the configurations of hydrogenation products and the reported mechanism of rhodium- and iridium-catalyzed asymmetric hydrogenation reactions,²¹ we proposed the transition model determining the enantioselectivity (Scheme 3). The ferrocenyl group of the ligand blocked one side of the complex and prevented the coordination of the substrate from this orientation. The substrate could only coordinate with rhodium from the opposing face of the catalyst. The repulsive steric interaction of the ligand with the larger sulfonyl group led to the coordination of the (*E*)-substrate with rhodium being more favorable from the *Re*-face than the *Si*-face of the α,β -unsaturated sulfones, predictably resulting in the (*R*)-product. By X-ray crystallographic analysis of the corresponding single-crystal structure or comparison with the sign of the specific rotation reported, the stereochemistry of some other products was determined. It was demonstrated to be in good agreement with the predicted stereochemical outcome of the hydrogenation products.

3. CONCLUSION

In conclusion, we have developed a highly efficient method for enantioselective synthesis of various optical active chiral sulfones. By employing a Rh-(*R,R*)-f-spiroPhos complex, a wide range of α,β -unsaturated sulfones including the (*E*)- and (*Z*)-isomers of 3,3-diaryl, cyclic, and 3-alkyl-3-aryl α,β -unsaturated sulfones as well as 2-alkyl-substituted benzo[*b*]thiophene 1,1-dioxides were successfully hydrogenated to the corresponding chiral sulfones with excellent enantioselectivities (up to 99.9% ee). Hydrogen pressure can be as low as 10 atm in certain cases with reaction completion in 1 h. Moreover, this method was also applied to the asymmetric synthesis of a spice flavor, (*S*)-(+)-*ar*-turmerone, an important synthetic intermediate of pharmaceuticals. We envision this method to allow for the broad screening of chiral sulfones for a wide variety of industries including pharmaceuticals, pesticides, or chiral polymers.

■ ASSOCIATED CONTENT**Supporting Information**

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.8b12657.

Experimental procedures, compound characterization data, and analysis of enantioselectivities of products (PDF)

Crystal parameters for compound 2l (CIF)

■ AUTHOR INFORMATION**Corresponding Author**

*ghhou@bnu.edu.cn

ORCID

Guofu Zi: 0000-0002-7455-460X

Guohua Hou: 0000-0002-3571-456X

Author Contributions

[†]Q.Y. and G.X. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the National Natural Science Foundation of China (Grant nos. 21672024, 21272026, and 21472013), Beijing Natural Science Foundation (2182025), and Beijing Municipal Commission of Education for generous financial support.

■ REFERENCES

(1) (a) Rupe, H.; Gassmann, A. Zur kenntnis des ar-turmerons aus dem cureuma-Öl. *Helv. Chim. Acta* **1936**, *19*, 569. (b) Serra, S. Preparation and use of enantioenriched 2-arylpropylsulfonylbenzene derivatives as valuable building blocks for the enantioselective synthesis of bisabolane sesquiterpenes. *Tetrahedron: Asymmetry* **2014**, *25*, 1561.

(2) Doswald, S.; Estermann, H.; Kupfer, E.; Stadler, H.; Walther, W.; Weisbrod, T.; Wirz, B.; Wostl, W. Large scale preparation of chiral building blocks for the P₃ site of renin inhibitors. *Bioorg. Med. Chem.* **1994**, *2*, 403.

(3) Liu, Y.; Qin, W.; Yan, H. Efficient enrichment of chiral β -sulfonyl ketones through asymmetric β -elimination. *Synlett* **2016**, *27*, 2756.

(4) Yamada, M.; Ichikawa, T.; Ii, M.; Itoh, K.; Tamura, N.; Kitazaki, T. Novel cyclohexene derivatives as anti-sepsis agents: Synthetic studies and inhibition of NO and cytokine production. *Bioorg. Med. Chem.* **2008**, *16*, 3941.

(5) (a) Trost, B. M. Organosulfons as synthetic building blocks. *Bull. Chem. Soc. Jpn.* **1988**, *61*, 107. (b) Nájera, C.; Yus, M. Desulfonylation reactions: recent developments. *Tetrahedron* **1999**, *55*, 10547. (c) Julia, M.; Paris, J.-M. Synthèses à l'aide de sulfones v⁽⁺⁾- methode de synthese generale de doubles liaisons. *Tetrahedron Lett.* **1973**, *14*, 4833.

(6) Choi, J.; Martín-Gago, P.; Fu, G. C. Stereoconvergent arylations and alkenylations of unactivated alkyl electrophiles: Catalytic enantioselective synthesis of secondary sulfonamides and sulfones. *J. Am. Chem. Soc.* **2014**, *136*, 12161.

(7) (a) Mauleon, P.; Carretero, J. C. Rhodium-catalyzed enantioselective conjugate addition of organoboronic acids to α,β -unsaturated sulfones. *Org. Lett.* **2004**, *6*, 3195. (b) Mauleon, P.; Carretero, J. C. Enantioselective construction of stereogenic quaternary centres via Rh-catalyzed asymmetric addition of alkenylboronic acids to α,β -unsaturated pyridylsulfones. *Chem. Commun.* **2005**, 4961. (c) Mauleon, P.; Alonso, I.; Rivero, M. R.; Carretero, J. C. Enantioselective synthesis of chiral sulfones by Rh-catalyzed asymmetric addition of boronic acids to α,β -unsaturated 2-pyridyl sulfones. *J. Org. Chem.* **2007**, *72*, 9924. (d) Mosse, S.; Alexakis, A. First organocatalyzed asymmetric michael addition of aldehydes to

vinyl sulfones. *Org. Lett.* **2005**, *7*, 4361. (e) Li, H. M.; Song, J.; Liu, X. F.; Deng, L. Catalytic enantioselective C-C bond forming conjugate additions with vinyl. *J. Am. Chem. Soc.* **2005**, *127*, 8948. (f) Liu, T. Y.; Long, J.; Li, B. J.; Jiang, L.; Li, R.; Wu, Y.; Ding, L. S.; Chen, Y. C. Enantioselective construction of quaternary carbon centre catalysed by bifunctional organocatalyst. *Org. Biomol. Chem.* **2006**, *4*, 2097. (g) Nielsen, M.; Jacobsen, C. B.; Holub, N.; Paixao, M. W.; Jørgensen, K. A. Asymmetric organocatalysis with sulfones. *Angew. Chem., Int. Ed.* **2010**, *49*, 2668.

(8) (a) Watanabe, Y.; Mase, N.; Furue, R.; Toru, T. Enantioselective allylation of the α -sulfonyl radical controlled by coordination of a chiral Lewis acid to an enantiotopic sulfonyl oxygen. *Tetrahedron Lett.* **2001**, *42*, 2981. (b) Sugimoto, H.; Nakamura, S.; Watanabe, Y.; Toru, T. Enantioselective hydrogen atom transfer to α -sulfonyl radicals controlled by selective coordination of a chiral Lewis acid to an enantiotopic sulfonyl oxygen. *Tetrahedron: Asymmetry* **2003**, *14*, 3043.

(9) Misun, M.; Pfaltz, A. Enantioselective reduction of electrophilic C = C Bonds with sodium tetrahydroborate and 'semicorrin' cobalt catalysts. *Helv. Chim. Acta* **1996**, *79*, 961.

(10) Llamas, T.; Arrayás, R. G.; Carretero, J. C. Catalytic asymmetric conjugate reduction of β,β -disubstituted α,β -unsaturated sulfones. *Angew. Chem., Int. Ed.* **2007**, *46*, 3329.

(11) Desrosiers, J.-N.; Charette, A. Catalytic enantioselective reduction of β,β -disubstituted vinyl phenyl sulfones by using bisphosphine monoxide ligands. *Angew. Chem., Int. Ed.* **2007**, *46*, 5955.

(12) (a) Ohkuma, T.; Kitamura, M.; Noyori, R. In *Catalytic Asymmetric Synthesis*, 2nd ed.; Ojima, I., Ed.; Wiley: New York, 2000; p 1. (b) Blaser, H.-U.; Spindler, F. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer: Berlin, 1999; p 247. (c) Tang, W.; Zhang, X. New chiral phosphorus ligands for enantioselective hydrogenation. *Chem. Rev.* **2003**, *103*, 3029. (d) Xie, J.-H.; Zhu, S.-F.; Zhou, Q.-L. Transition metal-catalyzed enantioselective hydrogenation of enamines and imines. *Chem. Rev.* **2011**, *111*, 1713. (e) Wang, D.-S.; Chen, Q.-A.; Lu, S.-M.; Zhou, Y.-G. Asymmetric hydrogenation of heteroarenes and arenes. *Chem. Rev.* **2012**, *112*, 2557. (f) Veredel, J. J.; Pàmies, O.; Diéguez, M.; Andersson, P. G. Asymmetric hydrogenation of olefins using chiral crabtree-type catalysts: scope and limitations. *Chem. Rev.* **2014**, *114*, 2130. (g) Xie, J.-H.; Zhou, Q.-L. New progress and prospects of transition metal-catalyzed asymmetric hydrogenation. *Huaxue Xuebao* **2012**, *70*, 1427. (h) Urban, S.; Beiring, B.; Ortega, N.; Paul, D.; Glorius, F. Asymmetric hydrogenation of thiophenes and benzothiophenes. *J. Am. Chem. Soc.* **2012**, *134*, 15241.

(13) (a) Zhou, T.; Peters, B.; Maldonado, M. F.; Govender, T.; Andersson, P. G. Enantioselective synthesis of chiral sulfones by Ir-catalyzed asymmetric hydrogenation: a facile approach to the preparation of chiral allylic and homoallylic compounds. *J. Am. Chem. Soc.* **2012**, *134*, 13592. (b) Peters, B. K.; Zhou, T.; Rujirawanich, J.; Cadu, A.; Singh, T.; Rabten, W.; Kerdphon, S.; Andersson, P. G. An enantioselective approach to the preparation of chiral sulfones by Ir-catalyzed asymmetric hydrogenation. *J. Am. Chem. Soc.* **2014**, *136*, 16557.

(14) Tosatti, P.; Pfaltz, A. Iridium-catalyzed asymmetric hydrogenation of benzo[b]thiophene 1,1-dioxides. *Angew. Chem., Int. Ed.* **2017**, *56*, 4579.

(15) Shi, L.; Wei, B.; Yin, X.; Xue, P.; Lv, H.; Zhang, X. Rh-catalyzed asymmetric hydrogenation of α -substituted vinyl sulfones: an efficient approach to chiral sulfones. *Org. Lett.* **2017**, *19*, 1024.

(16) Jiang, J.; Wang, Y.; Zhang, X. Rhodium-catalyzed asymmetric hydrogenation of β -acetylamino acrylsulfones: a practical approach to chiral β -amido sulfones. *ACS Catal.* **2014**, *4*, 1570.

(17) Paul, J. M.; Palmer, C. Asymmetric hydrogenation. U.S. Patent 6,274,758, 2001.

(18) (a) Tolstoy, P.; Engman, M.; Paptchikhine, A.; Bergquist, J.; Church, T. L.; Leung, A. W. M.; Andersson, P. G. Iridium-catalyzed asymmetric hydrogenation yielding chiral diarylmethines with weakly coordinating or noncoordinating substituents. *J. Am. Chem. Soc.* **2009**, *131*, 8855. (b) Mazuela, J.; Norrby, P.-O.; Andersson, P. G.; Pàmies,

O.; Diéguez, M. Pyranoside phosphite-oxazoline ligands for the highly versatile and enantioselective Ir-catalyzed hydrogenation of minimally functionalized olefins. a combined theoretical and experimental study. *J. Am. Chem. Soc.* **2011**, *133*, 13634. (c) Mazuela, J.; Verendel, J. J.; Coll, M.; Schäffner, B.; Börner, A.; Andersson, P. G.; Pamies, O.; Diéguez, M. Iridium phosphite-oxazoline catalysts for the highly enantioselective hydrogenation of terminal alkenes. *J. Am. Chem. Soc.* **2009**, *131*, 12344. (d) Boulton, L. T.; Lennon, I. C.; McCague, R. Asymmetric hydrogenation of a 4,4-diaryl-3-butenolate; a novel approach to sertraline. *Org. Biomol. Chem.* **2003**, *1*, 1094. (e) Wabnitz, T. C.; Rizzo, S.; Götte, C.; Buschauer, A.; Benincori, T.; Reiser, O. Enantioselective hydrogenation of diaryl-substituted α,β -unsaturated nitriles. *Tetrahedron Lett.* **2006**, *47*, 3733.

(19) (a) Yan, Q.; Liu, M.; Kong, D.; Zi, G.; Hou, G. Highly efficient iridium-catalyzed asymmetric hydrogenation of β -acylamino nitroolefins. *Chem. Commun.* **2014**, *50*, 12870. (b) Yan, Q.; Kong, D.; Li, M.; Hou, G.; Zi, G. Highly efficient Rh-catalyzed asymmetric hydrogenation of α,β -unsaturated nitriles. *J. Am. Chem. Soc.* **2015**, *137*, 10177. (c) Liu, M.; Kong, D.; Li, M.; Zi, G.; Hou, G. Iridium-catalyzed enantioselective hydrogenation of β,β -disubstituted nitroalkenes. *Adv. Synth. Catal.* **2015**, *357*, 3875. (d) Kong, D.; Li, M.; Wang, R.; Zi, G.; Hou, G. Asymmetric hydrogenation of β -aryloxy/alkoxy cinnamic nitriles and esters. *Org. Lett.* **2016**, *18*, 4916. (e) Kong, D.; Li, M.; Zi, G.; Hou, G.; He, Y. Enantioselective hydrogenation of diarylmethanimines for synthesis of chiral diarylmethylamines. *J. Org. Chem.* **2016**, *81*, 6640. (f) Yan, Q.; Kong, D.; Zhao, W.; Zi, G.; Hou, G. Enantioselective hydrogenation of β,β -disubstituted unsaturated carboxylic acids under base-free conditions. *J. Org. Chem.* **2016**, *81*, 2070. (g) Zhang, Y.; Yan, Q.; Zi, G.; Hou, G. Enantioselective direct synthesis of free cyclic amines via intramolecular reductive amination. *Org. Lett.* **2017**, *19*, 4215. (h) Li, M.; Kong, D.; Zi, G.; Hou, G. Rh-catalyzed asymmetric hydrogenation of 1,2-dicyanoalkenes. *J. Org. Chem.* **2017**, *82*, 680.

(20) (R)-JosiPhos: 30% conversion, 42% ee; (S,S)-Me-Duphos: 2.3% conversion; (R,R)-Ph-BPE: 2.2% conversion.

(21) (a) Gridnev, I. D.; Higashi, N.; Asakura, K.; Imamoto, T. Mechanism of asymmetric hydrogenation catalyzed by a rhodium complex of (S,S)-1,2-bis(*tert*-butylmethylphosphino)ethane. Dihydride mechanism of asymmetric hydrogenation. *J. Am. Chem. Soc.* **2000**, *122*, 7183. (b) Gridnev, I. D.; Yasutake, M.; Higashi, N.; Imamoto, T. Asymmetric hydrogenation of enamides with Rh-BisP* and Rh-MiniPHOS catalysts. Scope, limitations, and mechanism. *J. Am. Chem. Soc.* **2001**, *123*, 5268. (c) Li, J.-Q.; Quan, X.; Andersson, P. G. Highly enantioselective iridium-catalyzed hydrogenation of α,β -unsaturated esters. *Chem. - Eur. J.* **2012**, *18*, 10609.