

# Rare-Earth-Catalyzed Selective 1,4-Hydrosilylation of Branched 1,3-Enynes Giving Tetrasubstituted Silyllallenes

Wufeng Chen, Chunhui Jiang, Jianying Zhang, Jiaqi Xu, Lin Xu, Xiufang Xu, Jianfeng Li,\* and Chunming Cui\*

Cite This: *J. Am. Chem. Soc.* 2021, 143, 12913–12918

Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information

**ABSTRACT:** Allenes are versatile synthons in organic synthesis and medicinal chemistry because of their diverse reactivities. Catalytic 1,4-hydrosilylation of 1,3-enynes may present the straightforward strategy for synthesis of silyllallenes. However, the transition-metal-catalyzed reaction has not been successful due to poor selectivity and very limited substrate scopes. We report here the efficient and selective 1,4-hydrosilylation of branched 1,3-enynes enabled by the ene-diamido rare-earth ate catalysts using both alkyl and aryl hydrosilanes, leading to the exclusive formation of tetrasubstituted silyllallenes. Deuteration reaction, kinetic study, and DFT calculations were conducted to investigate the possible mechanism, revealing crucial roles of high Lewis acidity, large ionic radius, and ate structure of the rare-earth catalysts.

Metal-catalyzed hydrosilylation of unsaturated organic compounds, especially alkenes and alkynes, represents the most efficient and straightforward protocol for the synthesis of various silyl-functionalized organic molecules, which played important roles in organic and materials chemistry.<sup>1</sup> 1,3-Enynes, readily available conjugated organic compounds containing both alkene and alkyne moieties, are basic building blocks in synthetic chemistry.<sup>2</sup> However, catalytic hydrosilylation of 1,3-enynes confronts the issues of regio- and stereoselective control because there exist three mainly competing pathways, namely, 1,2-, 4,3-, and 1,4-hydrosilylation (Scheme 1).<sup>3</sup> As a result, only a few cases of transition-metal-catalyzed 4,3-hydrosilylation of 1,3-enynes for the synthesis of silyldienes have been reported,<sup>4</sup> while catalytic 1,4-hydrosilylation of 1,3-enynes to silyllallenes has been met with very little success. There were only a couple of Pt- and

Pd-catalyzed asymmetric 1,4-hydrosilylations of 1,3-enynes that have been reported.<sup>3a,b,5</sup> However, these precious-metal-catalyzed reactions suffered from very limited substrate scopes. Because allenenes feature unique structures and diverse reactivities and are valuable multifunctional synthons in synthetic and medicinal chemistry,<sup>6</sup> the development of efficient and selective catalysts for 1,4-hydrosilylation of enynes is highly desired.


Rare-earth-catalyzed hydrosilylation of alkenes and alkynes exhibits unique reactivity and selectivity because of their high Lewis acidity and large ionic radii.<sup>7</sup> In particular, we have recently shown that ene-diamido rare-earth (RE) complexes enabled highly regioselective hydrosilylation of internal alkenes and alkynes.<sup>8</sup> Stimulated by the high reactivity of these rare-earth complexes, we are interested in the exploration of rare-earth-catalyzed hydrosilylation of 1,3-enynes with the expectation to realize some novel and selective transformations for the synthesis of valuable synthons and organosilanes.

Herein, we report the employment of ene-diamido rare-earth ate complexes for the successful catalytic 1,4-hydrosilylation of branched 1,3-enynes, leading to the formation of tetrasubstituted silyllallenes (Scheme 1). Mechanistic studies and DFT calculations indicated that the high Lewis acidity, large ionic radius, and unique ate structure are key factors for the unprecedented transformation.


Catalytic hydrosilylation of 2-methyl-4-trimethylsilyl-1-butene-3-yne (7a) with  $n\text{C}_6\text{H}_{13}\text{SiH}_3$  (6a) was selected as model reaction (Table 1). Initially, we investigated the ene-diamido

## Scheme 1. Catalytic Hydrosilylation of 1,3-Enynes

**Previous Work:** Transition-Metal-Catalyzed Hydrosilylation of Enynes




**This Work:** Rare-Earth-Catalyzed 1,4-Hydrosilylation of Branched Enynes



Received: May 5, 2021

Published: August 13, 2021



Table 1. Reaction Condition Optimization<sup>a</sup>

entry	cat	solvent	time (h)	T (°C)	yield (%) <sup>b</sup>
1	YC	C <sub>6</sub> D <sub>6</sub>	12	80	<5
2	1	C <sub>6</sub> D <sub>6</sub>	12	80	<5
3	3	C <sub>6</sub> D <sub>6</sub>	12	80	<5
4	2	C <sub>6</sub> D <sub>6</sub>	4	rt	74
5	4	C <sub>6</sub> D <sub>6</sub>	4	rt	68
6	5	C <sub>6</sub> D <sub>6</sub>	4	rt	21
7	2	C <sub>6</sub> D <sub>6</sub>	8	rt	>99
8	KN <sup>c</sup>	C <sub>6</sub> D <sub>6</sub>	24	80	29
9	2	toluene	8	rt	>99
10	2	<i>n</i> -hexane	8	rt	52
11	2	THF	24	80	<5


<sup>a</sup>*n*C<sub>6</sub>H<sub>13</sub>SiH<sub>3</sub> (6a) (0.44 mmol), 2-methyl-4-trimethylsilyl-1-butene-3-yne (7a) (0.4 mmol), catalyst 2 (0.02 mmol, 5 mol %), 0.6 mL of solvent. <sup>b</sup>Yields of 8a were determined by <sup>1</sup>H NMR and GC-MS with the crude mixture. <sup>c</sup>KN = KN(SiMe<sub>3</sub>)<sub>2</sub>.

rare-earth alkyl and amide complexes LY(CH<sub>2</sub>SiMe<sub>3</sub>)(THF)<sub>2</sub> (YC, L = ArNC(Me)C(Me)NAr, Ar = 2,6-*i*Pr<sub>2</sub>C<sub>6</sub>H<sub>3</sub>, entry 1), LSm(CH<sub>2</sub>SiMe<sub>3</sub>)(THF)<sub>2</sub> (1, entry 2),<sup>8b</sup> and LLa(N(SiMe<sub>3</sub>)<sub>2</sub>)(THF) (3, entry 3).<sup>8c</sup> However, only very low conversions (<5% yield of geminal disilyllallene 8a) have been observed in 12 h at 80 °C with these neutral catalysts. Prompted by our recent successes in the hydrosilylation of internal alkynes with rare-earth ate complexes, the ene-diamido lanthanum bis-(amido) ate complex K[LLa(N(SiMe<sub>3</sub>)<sub>2</sub>)<sub>2</sub>] 2 (entry 4) was employed. Indeed, 2 enabled highly selective hydrosilylation of 7a with 6a to give 1,4-hydrosilylation product 8a in high yield (74%) in 4 h at room temperature without the contamination of the other hydrosilylation products. The samarium and yttrium ate complexes K[LSm(N(SiMe<sub>3</sub>)<sub>2</sub>)<sub>2</sub>] (4)<sup>8c</sup> and K[LY(N(SiMe<sub>3</sub>)<sub>2</sub>)<sub>2</sub>] (5, see the Supporting Information (SI)) were also examined (entries 5 and 6). Under same reaction conditions, the samarium ate complex 4 gave 8a in 68% yield whereas the yttrium complex 5 only produced 8a in 21% yield. This activity trend appeared to be closely correlated with the ionic radii of the elements: the larger the ionic radius is, the higher the activity (La<sup>3+</sup> > Sm<sup>3+</sup> > Y<sup>3+</sup>). When the reaction time was extended to 8 h (entry 7), the lanthanum ate complex 2 gave 8a almost quantitatively. KN(SiMe<sub>3</sub>)<sub>2</sub> (KN, entry 8) is also active but only gave 8a in 29% yield at 80 °C in 24 h, and the reaction did not produce any more product after 24 h. These results indicated that the combination of a large rare-earth ion with KN(SiMe<sub>3</sub>)<sub>2</sub> forming ate structure is crucial for the effective catalytic 1,4-hydrosilylation of enynes.

Toluene (entry 9) and C<sub>6</sub>D<sub>6</sub> are the superior solvents for the hydrosilylation of the enyne, whereas the reaction in *n*-hexane (entry 10) only resulted in the 1,4-hydrosilylation product in 52% yield probably because of the poor solubility of 2 in *n*-

hexane. In sharp contrast, almost no reaction was observed in THF (entry 11) probably due to the interactions of the solvent molecules with the lanthanum ion, which suppresses the coordination–insertion process.

Under the optimized conditions (Table 1, entry 9), hydrosilylation of a range of branched 1,3-enynes 7a–r with alkyl-substituted hydrosilanes *n*C<sub>6</sub>H<sub>13</sub>SiH<sub>3</sub> 6a and CySiH<sub>3</sub> 6b was examined (Table 2). All of the branched enynes

Table 2. Hydrosilylation with Alkyl Hydrosilanes<sup>a</sup>

<sup>a</sup>RSiH<sub>3</sub> 6a and 6b (R = *n*C<sub>6</sub>H<sub>13</sub>, R = cyclohexyl, 0.44 mmol), enynes 7a–r (0.4 mmol), catalyst 2 (0.02 mmol, 5 mol %), rt and 1 mL of toluene. Isolated yield. <sup>b</sup>60 °C.


underwent selective 1,4-hydrosilylation to afford tetrasubstituted silyllallenes 8a–r in excellent yields (83–94%), while the other hydrosilylation products were not observable. When the internal substituent R<sup>2</sup> on the alkene moiety is alkyl, the catalytic reaction could be applied to enynes containing various silyl-substituted alkyne moieties (R<sup>1</sup>) to yield geminal disilyllallenes (8a–g); when the internal substituent R<sup>2</sup> is silyl, the reaction is viable for enynes containing alkyl-substituted alkyne moieties (R<sup>1</sup>) to produce 1,3-disilyllallenes (8h–q). It is noteworthy that the 1,4-hydrosilylation underwent smoothly in good yields for the enynes with sterically demanding groups (86–87% yields of 8c, 8d, and 8m). The

hydrosilylation of enynes with siloxyl and N-heterocyclic substituents (**8p** and **8q**) could also achieve good yields in 16 and 8 h. The enynes with chloro and hydroxyl groups were not compatible with this reaction.

The substituent effects of 1,3-enynes on the hydrosilylation selectivity have been further examined with alkyl substituted enynes and terminal enynes. 1,3-Enynes with *t*-butyl and cyclohexyl groups could also be successfully hydrosilylated to give 1,4-hydrosilylation products (**8r** and **8s**, 91 and 92% yields) in 12 and 6 h. However, *n*-hexyl ( $R^1$ ) substituted 1,3-enynes ( $R^2 = \text{H, Me}$ ) exhibited poor regioselectivity with the formation of 1,4- and 4,3-hydrosilylation products (Table S1 in the SI). It is very likely that the steric effects of  $R^1$  group could efficiently inhibit the 4,3-insertion but favor 1,4-selectivity.<sup>10</sup> It is noted that **8n** was generated selectively from enynes with  $R^1 = n$ -hexyl and  $R^2 =$  silyl groups because of the inductive effects of the silyl group on the alkene moiety, which could result in high electron density on the silyl carbon atom and facilitate the 1,2-insertion.<sup>8c,11</sup> However, with 1,3-enynes of  $R^1 =$  aryl and  $R^2 =$  silyl groups, the reactions only yielded polymeric materials and hydrosilylation products could not be observed.<sup>8a,12</sup>

Aryl-substituted hydrosilanes could also be applied to the catalytic reaction (Table 3) with satisfactory yields (81–90%)

Table 3. Hydrosilylation with Aryl Hydrosilanes<sup>a</sup>




<sup>a</sup>ArSiH<sub>3</sub> **6c-j** (0.44 mmol), 1,3-enyne **7a**, **7g**, and **7m** (0.4 mmol), catalyst **2** (0.02 mmol, 5 mol %), rt and 1 mL of toluene. Isolated yield. <sup>b</sup>60 °C.

and excellent regioselectivity. The catalytic protocol is tolerated to some functional groups (chloro **8v** and alkoxy **8w**). In addition, the bis(allene) **8y** can be obtained. It is noted that the substituents of hydrosilanes have significant effect on the catalytic activity: aryl hydrosilanes (phenyl **8t**) are more active than alkyl hydrosilanes (*n*-hexyl **8a** and cyclohexyl **8b**), and bulky hydrosilanes are less active (naphthyl **8x**). To our delight, the 1,4-hydrosilylation with the secondary silane Ph<sub>2</sub>SiH<sub>2</sub> proceeded successfully to give the expected product **9** in 84% yield. The hydrosilylation with PhSiD<sub>3</sub> yielded the deuteration 1,3-disilyllallenes (**10**) exclusively (90% yield),


indicating the 1,4-hydrosilylation by D and RSiD<sub>2</sub> groups, which was strongly supported by the <sup>1</sup>H, <sup>2</sup>H, and <sup>13</sup>C NMR spectra. The synthetic potentials of silyllallenes was demonstrated by the desilylation of **8o** with TBAF (Scheme 2), which afforded 1,3-disubstituted allene **11** in 90% yield.<sup>13</sup>

Scheme 2. Derivatization of Silyllallene



The reaction of lanthanum ate complex **2** with silane **6** was reported to yield lanthanum hydride  $\text{K}[\text{L}((\text{Me}_3\text{Si})_2\text{N})\text{La}(\mu\text{-H})_2\text{La}(\text{H})(\text{THF})\text{L})\text{K}]$ .<sup>8c</sup> In contrast, the neutral complex **3** is inactive to the hydrosilane. These results indicated that the negative charge in anionic  $[\text{LLa}(\text{N}(\text{SiMe}_3)_2)_2]^-$  of ate complex **2** could facilitate the  $\sigma$ -bond metathesis of **2** with hydrosilane to form active rare-earth hydride intermediate (Figure 1a).<sup>8,14,15</sup> The catalytic cycle may initiate with the selective 1,2-insertion of enyne **7** into the lanthanum hydride **A**, giving the  $\eta^3$ -propargyl/allenyl lanthanum intermediate **B**, similar to the Cu and Mg hydride-catalyzed 1,4-hydroelementation of 1,3-enynes.<sup>10</sup> After that,  $\sigma$ -bond metathesis of **B** with hydrosilane **6** led to the exclusive formation of silyllallene **8** and regeneration of the active hydride **A**.

To gain further support for the proposed mechanism, DFT calculations were carried out. First, the LUMO of the 1,3-enyne **7g** was calculated using frontier orbital population analysis (Figure S1 in the SI). The coefficients at the C1 and C2 (0.40, 0.31) atoms of the alkene moiety are larger than those at the C4 and C3 (0.27, 0.13) atoms of the alkyne moiety, which explains, to a certain degree, the preference for initial 1,2-insertion rather than 4,3-insertion.<sup>16</sup> To further understand the mechanism, the calculations were also performed on the 1,4-hydrosilylation of 1,3-enyne **7a** with MeSiH<sub>3</sub> using  $[\text{LLa}(\text{H})(\text{N}(\text{SiMe}_3)_2)_2]^-$  as the catalyst (Figure 1b). It is predicted that the hydrosilylation was initiated by the facile 1,2-insertion of an enyne via the four-centered TS1 (Figure 1c) with the energy barrier of 21.1 kcal/mol, affording the  $\eta^3$ -propargyl/allenyl lanthanum intermediate **B**. The 4,3-insertion showed higher activation energy ( $\Delta\Delta G^\ddagger = 0.4$  kcal/mol) than that of 1,2-insertion because of the strong steric repulsion between substituents on the internal alkyne moiety and bulky ene-diamido ligand in TS1' (Figure S2 in the SI). Meanwhile, the calculations suggest that propargyl/allenyl intermediate **B** from 1,2-insertion is more stable ( $\Delta\Delta G = -4.7$  kcal/mol) than dienyl intermediate **B'** from 4,3-insertion. It is worth noting that the  $\eta^1$ -propargyl lanthanum species has not been predicted on the basis of 1,2-insertion, which is probably caused by the large lanthanum ion that could favor the  $\eta^3$ -fashion.<sup>17</sup> In fact, the short distance of La–C4 (2.67 Å) compared to that of La–C2 (3.37 Å), along with the almost linear C2–C3–C4 angle of 171.9°, suggests that the geometry of intermediate **B** lies between  $\eta^3$ -propargyl/allenyl and  $\eta^1$ -allenyl structures. Finally, intermediate **B** underwent  $\sigma$ -bond metathesis with hydrosilane via the TS2 to give the silyllallene with the regeneration of the hydride intermediate. This  $\sigma$ -bond metathesis step requires a higher energy barrier (24.1 kcal/mol) and thus becomes the rate-limiting step of the overall reaction.



**Figure 1.** (a) Proposed mechanism. (b) Gibbs free-energy profile. (c) Calculated structures of TS1, B, and TS2.

Kinetic studies on 1,4-hydrosilylation were carried out (Table S2 and Figure S3 in the SI) which indicated that the rate law was of first order dependence on the hydrosilane and enyne.<sup>8c</sup> The results suggested that the reaction rate was influenced by both substrates. It is probably attributed to the steric hindrance of the branched 1,3-enynes, which leads to the intermediate B with bulky  $\eta^3$ -propargyl/allenyl features having a relatively slow  $\sigma$ -bond metathesis reaction with hydrosilane.

In summary, we have disclosed that the lanthanum ate complex 2 enabled highly efficient and selective 1,4-hydrosilylation of branched 1,3-enynes with various primary and secondary hydrosilanes to yield tetrasubstituted silyllallenes. The high activity for sterically demanding branched enynes could be attributed to the high Lewis acidity and large ionic radius of lanthanum ion as well as the highly reactive ate hydride intermediate. Further applications of ene-diamido rare-earth ate catalysts in hydroelementation of bulky unsaturated molecules are in progress.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.1c04689>.

Experimental procedures, DFT calculations, and spectroscopic data (PDF)

## AUTHOR INFORMATION

### Corresponding Authors

Jianfeng Li – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China; [orcid.org/0000-0003-0996-1679](https://orcid.org/0000-0003-0996-1679); Email: [lijf@nankai.edu.cn](mailto:lijf@nankai.edu.cn)

Chunming Cui – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China; [orcid.org/0000-0002-1493-7596](https://orcid.org/0000-0002-1493-7596); Email: [cmcui@nankai.edu.cn](mailto:cmcui@nankai.edu.cn)

### Authors

Wufeng Chen – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China

Chunhui Jiang – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China

Jiaying Zhang – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China; [orcid.org/0000-0001-6119-2497](https://orcid.org/0000-0001-6119-2497)

Jiaqi Xu – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China

Lin Xu – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China

Xiufang Xu – State Key Laboratory of Elemento-Organic Chemistry and College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China; [orcid.org/0000-0002-3510-3267](https://orcid.org/0000-0002-3510-3267)

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jacs.1c04689>

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We are grateful to the National Natural Science Foundation of China (Grant No. 21890722, 22071123 and 21632006) for financial support. Dedicated to the 100th Anniversary of Chemistry at Nankai University.

## REFERENCES

- (1) (a) Marciniak, B.; Maciejewski, H.; Pietraszuk, C.; Pawluć, P. In *Hydrosilylation: A Comprehensive Review on Recent Advances*; Marciniak, B., Ed.; Springer: Berlin, 2009; Vol. 1, pp 3–154. (b) Trost, B. M.; Ball, Z. T. Addition of Metalloid Hydrides to Alkynes: Hydrometallation with Boron, Silicon, and Tin. *Synthesis* **2005**, 2005, 853–887. (c) Troegel, D.; Stohrer, J. Recent advances and actual challenges in late transition metal catalyzed hydrosilylation of olefins from an industrial point of view. *Coord. Chem. Rev.* **2011**, 255, 1440–1459. (d) Lim, D. S. W.; Anderson, E. A. Synthesis of Vinylsilanes. *Synthesis* **2012**, 44, 983–1010. (e) Nakajima, Y.; Shimada, S. Hydrosilylation reaction of olefins: recent advances and perspectives. *RSC Adv.* **2015**, 5, 20603–20616. (f) Sun, J.; Deng, L.



Cobalt Complex-Catalyzed Hydrosilylation of Alkenes and Alkynes. *ACS Catal.* **2016**, *6*, 290–300. (g) Du, X.; Huang, Z. Advances in Base-Metal-Catalyzed Alkene Hydrosilylation. *ACS Catal.* **2017**, *7*, 1227–1243. (h) Zaranek, M.; Pawluc, P. Markovnikov Hydro-silylation of Alkenes: How an Oddity Becomes the Goal. *ACS Catal.* **2018**, *8*, 9865–9876.

(2) (a) Wessig, P.; Müller, G. The Dehydro-Diels–Alder Reaction. *Chem. Rev.* **2008**, *108*, 2051–2063. (b) Dherbassy, Q.; Manna, S.; Talbot, F. J. T.; Prasitwatcharakorn, W.; Perry, G. J. P.; Procter, D. J. Copper-catalyzed functionalization of enynes. *Chem. Sci.* **2020**, *11*, 11380–11393.

(3) (a) Kusumoto, T.; Hiyama, T. Hydrosilylation of 1,4-bis(trimethylsilyl)-1,3-butadiyne. *Chem. Lett.* **1985**, *14*, 1405–1408. (b) Kusumoto, T.; Ando, K.; Hiyama, T. Hydrosilylation of 1,4-bis(trimethylsilyl)butadiyne and silyl-substituted butenyne. *Bull. Chem. Soc. Jpn.* **1992**, *65*, 1280–1290. (c) Maruyama, Y.; Yoshiuchi, K.; Ozawa, F.; Wakatsuki, Y. Regio- and stereoselective hydrosilylation of 1,4-bis(trimethylsilyl)-3-buten-1-yne. *Chem. Lett.* **1997**, *26*, 623–624. (d) Maruyama, Y.; Yoshiuchi, K.; Ozawa, F. Hydrosilylation of 1,4-bis(trimethylsilyl)-1-buten-3-yne using late transition metal hydrides as catalyst precursors. *J. Organomet. Chem.* **2000**, *609*, 130–136. (e) Choudhury, P. P.; Junker, C. S.; Pidaparthy, R. R.; Welker, M. E. Syntheses of 2-silicon-substituted 1,3-dienes. *J. Organomet. Chem.* **2014**, *754*, 88–93.

(4) (a) Micoine, K.; Furstner, A. Concise Total Synthesis of the Potent Translation and Cell Migration Inhibitor Lactimidomycin. *J. Am. Chem. Soc.* **2010**, *132*, 14064–14066. (b) Gallenkamp, D.; Furstner, A. Stereoselective Synthesis of E,Z-Configured 1,3-Dienes by Ring-Closing Metathesis. Application to the Total Synthesis of Lactimidomycin. *J. Am. Chem. Soc.* **2011**, *133*, 9232–9235. (c) Zhou, H.; Moberg, C. Regio- and Stereoselective Hydrosilylation of 1,3-Enynes Catalyzed by Palladium. *Org. Lett.* **2013**, *15*, 1444–1447. (d) Trost, B. M.; Ball, Z. T.; Jöge, T. Regioselective Hydrosilylation of Propargylic Alcohols: An Aldol Surrogate. *Angew. Chem., Int. Ed.* **2003**, *42*, 3415–3418. (e) Bergueiro, J.; Montenegro, J.; Cambeiro, F.; Saa, C.; López, S. Cross-Coupling Reactions of Organosilicon Compounds in the Stereocontrolled Synthesis of Retinoids. *Chem. - Eur. J.* **2012**, *18*, 4401–4410. (f) Bergueiro, J.; Montenegro, J.; Saa, C.; López, S. Synthesis of 11-cis-Retinoids by Hydrosilylation-Protodesilylation of an 11,12-Didehydro Precursor: Easy Access to 11- and 12-Mono- and 11,12-Dideuteroretinoids. *Chem. - Eur. J.* **2012**, *18*, 14100–14107. (g) Walkowiak, J.; Salamon, K.; Franczyk, A.; Stefanowska, K.; Szyling, J.; Kownacki, I. Pt-Catalyzed Hydrosilylation of 1,3-Diynes with Triorganosilanes: Regio- and Stereoselective Synthesis of Mono- or Bis-silylated Adducts. *J. Org. Chem.* **2019**, *84*, 2358–2365. (h) Guo, Z.; Wen, H.; Liu, G.; Huang, Z. Iron-Catalyzed Regio- and Stereoselective Hydrosilylation of 1,3-Enynes To Access 1,3-Dienylsilanes. *Org. Lett.* **2021**, *23*, 2375–2379.

(5) (a) Han, J. W.; Tokunaga, N.; Hayashi, T. Palladium-Catalyzed Asymmetric Hydrosilylation of 4-Substituted 1-Buten-3-yne. Catalytic Asymmetric Synthesis of Axially Chiral Allenylsilanes. *J. Am. Chem. Soc.* **2001**, *123*, 12915–12916. (b) Ogasawara, M.; Ito, A.; Yoshida, K.; Hayashi, T. Synthesis of 2,5-Bis(binaphthyl)phospholes and Phosphametalloocene Derivatives and Their Application in Palladium-Catalyzed Asymmetric Hydrosilylation. *Organometallics* **2006**, *25*, 2715–2718.

(6) (a) Hopf, H. In *Modern Allene Chemistry*; Krause, N., Hashimi, A. S. K., Eds.; Wiley-VCH: Weinheim, 2004; pp 185–241. (b) Zimmer, R.; Dinesh, C. U.; Nandan, E.; Khan, F. A. Palladium-Catalyzed Reactions of Allenes. *Chem. Rev.* **2000**, *100*, 3067–3126. (c) Ma, S. Transition Metal-Catalyzed/Mediated Reaction of Allenes with a Nucleophilic Functionality Connected to the  $\alpha$ -Carbon Atom. *Acc. Chem. Res.* **2003**, *36*, 701–712. (d) Sydnes, L. K. Allenes from Cyclopropanes and Their Use in Organic Synthesis Recent Developments. *Chem. Rev.* **2003**, *103*, 1133–1150. (e) Hoffmann-Röder, A.; Krause, N. Synthesis and Properties of Allenic Natural Products and Pharmaceuticals. *Angew. Chem., Int. Ed.* **2004**, *43*, 1196–1216. (f) Hammond, G. B. Functionalized Fluorinated Allenes. *ACS Symp. Ser.* **2005**, *911*, 204–215. (g) Ma,

S. Some Typical Advances in the Synthetic Applications of Allenes. *Chem. Rev.* **2005**, *105*, 2829–2872. (h) Ma, S. Electrophilic Addition and Cyclization Reactions of Allenes. *Acc. Chem. Res.* **2009**, *42*, 1679–1688. (i) Rivera-Fuentes, P.; Diederich, F. Allenes in Molecular Materials. *Angew. Chem., Int. Ed.* **2012**, *51*, 2818–2828. (j) Brasholz, M.; Reissig, H.-U.; Zimmer, R. Sugars, Alkaloids, and Heteroaromatics: Exploring Heterocyclic Chemistry with Alkoxyallenes. *Acc. Chem. Res.* **2009**, *42*, 45–56. (k) Yu, S.; Ma, S. Allenes in Catalytic Asymmetric Synthesis and Natural Product Syntheses. *Angew. Chem., Int. Ed.* **2012**, *51*, 3074–3112. (l) Lu, T.; Lu, Z.; Ma, Z.-X.; Zhang, Y.; Hsung, R. P. Allenamides: A Powerful and Versatile Building Block in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 4862–4904. (m) Allen, A. D.; Tidwell, T. T. Ketenes and Other Cumulenes as Reactive Intermediates. *Chem. Rev.* **2013**, *113*, 7287–7342. (n) Ye, J.; Ma, S. Palladium-Catalyzed Cyclization Reactions of Allenes in the Presence of Unsaturated Carbon–Carbon Bonds. *Acc. Chem. Res.* **2014**, *47*, 989–1000. (o) López, F.; Mascareñas, J. L. [4 + 2] and [4 + 3] catalytic cycloadditions of allenenes. *Chem. Soc. Rev.* **2014**, *43*, 2904–2915. (p) Alcaide, B.; Almendros, P.; Aragoncillo, C. Cyclization reactions of bis(allenes) for the synthesis of polycarbo(hetero)cycles. *Chem. Soc. Rev.* **2014**, *43*, 3106–3135. (q) Muñoz, M. P. Silver and platinum-catalysed addition of O–H and N–H bonds to allenenes. *Chem. Soc. Rev.* **2014**, *43*, 3164–3183. (r) Neff, R. K.; Frantz, D. E. Recent Advances in the Catalytic Syntheses of Allenes: A Critical Assessment. *ACS Catal.* **2014**, *4*, 519–528. (s) Holmes, M.; Schwartz, L. A.; Krische, M. J. Intermolecular Metal-Catalyzed Reductive Coupling of Dienes, Allenes, and Enynes with Carbonyl Compounds and Imines. *Chem. Rev.* **2018**, *118*, 6026–6052.

(7) (a) Anwender, R. In *Principles in Organolanthanide Chemistry*; Springer: Berlin/Heidelberg, 1999; pp 3–23. (b) Reznichenko, A. L.; Hultsch, K. C. In *Molecular Catalysis of Rare-Earth Elements*; Roesky, P. W., Ed.; Springer: Berlin, 2010; Vol. 137, pp 2–12. (c) Molander, G. A.; Romero, J. A. C. Lanthanocene Catalysts in Selective Organic Synthesis. *Chem. Rev.* **2002**, *102*, 2161–2186. (d) Liu, D.; Liu, B.; Pan, Z.; Li, J.; Cui, C. Rare-earth metal catalysts for alkene hydrosilylation. *Sci. China: Chem.* **2019**, *62*, 571–682.

(8) (a) Li, J.; Zhao, C.; Liu, J.; Huang, H.; Wang, F.; Xu, X.; Cui, C. Activation of Ene-Diamido Samarium Methoxide with Hydrosilane for Selectively Catalytic Hydrosilylation of Alkenes and Polymerization of Styrene: an Experimental and Theoretical Mechanistic Study. *Inorg. Chem.* **2016**, *55*, 9105–9111. (b) Liu, J.; Chen, W.; Li, J.; Cui, C. Rare-Earth-Catalyzed Regioselective Hydrosilylation of Aryl-Substituted Internal Alkenes. *ACS Catal.* **2018**, *8*, 2230–2235. (c) Chen, W.; Song, H.; Li, J.; Cui, C. Catalytic Selective Dihydrosilylation of Internal Alkynes Enabled by Rare-Earth Ate Complex. *Angew. Chem., Int. Ed.* **2020**, *59*, 2365–2369. (d) Chen, W.; Liu, X.; Kong, J.; Li, J.; Chen, Y.; Cui, C. Rare-Earth-Catalyzed Selective Synthesis of Linear Hydridopolycarbosilanes and Their Functionalization. *Macromolecules* **2021**, *54*, 673–678.

(9) Kissel, A. A.; Mahrova, T. V.; Lyubov, D. M.; Cherkasov, A. V.; Fukin, G. K.; Trifonov, A. A.; Del Rosal, I.; Maron, L. Metallacyclic yttrium alkyl and hydrido complexes: synthesis, structures and catalytic activity in intermolecular olefin hydrophosphination and hydroamination. *Dalton Trans.* **2015**, *44*, 12137–12148.

(10) (a) Huang, Y.; del Pozo, J.; Torker, S.; Hoveyda, A. H. Enantioselective Synthesis of Trisubstituted Allenyl–B(pin) Compounds by Phosphine–Cu-Catalyzed 1,3-Enyne Hydroboration. Insights Regarding Stereochemical Integrity of Cu–Allenyl Intermediates. *J. Am. Chem. Soc.* **2018**, *140*, 2643–2655. (b) Wang, B.; Li, Y.; Pang, J. H.; Watanabe, K.; Takita, R.; Chiba, S. Hydro-magnesiation of 1,3-Enynes by Magnesium Hydride for Synthesis of Tri- and Tetra-substituted Allenes. *Angew. Chem., Int. Ed.* **2021**, *60*, 217–221.

(11) (a) Snider, B. B.; Karras, M.; Conn, R. S. E. Nickel-catalyzed addition of Grignard reagents to silylacetylenes. Synthesis of tetrasubstituted alkenes. *J. Am. Chem. Soc.* **1978**, *100*, 4624–4626. (b) Gao, Y.; Sato, F. On the mechanism of titanocenedichloride-catalysed hydromagnesiation of alkynes with alkyl Grignard reagents. *J. Chem. Soc., Chem. Commun.* **1995**, 659–660. (c) Shirakawa, E.;

Yamagami, T.; Kimura, T.; Yamaguchi, S.; Hayashi, T. Arylmagnesium of Alkynes Catalyzed Cooperatively by Iron and Copper Complexes. *J. Am. Chem. Soc.* **2005**, *127*, 17164–17165. (d) Tan, B. H.; Dong, J.; Yoshikai, N. Cobalt-catalyzed addition of arylzinc reagents to alkynes to form ortho-alkenylarylzinc species through 1,4-cobalt migration. *Angew. Chem., Int. Ed.* **2012**, *51*, 9610–9614. (e) Lee, Y. H.; Morandi, B. Palladium-Catalyzed Intermolecular Aryliodination of Internal Alkynes. *Angew. Chem., Int. Ed.* **2019**, *58*, 6444–6448. (f) Pan, Z.; Gao, D.; Zhang, C.; Guo, L.; Li, J.; Cui, C. Synthesis and Reactivity of N-heterocyclic Carbene Stabilized Lanthanide(II) Bis(amido) Complexes. *Organometallics* **2021**, *40*, 1728–1734.

(12) Fu, P.-F.; Marks, T. J. Silanes as Chain Transfer Agents in Metallocene-Mediated Olefin Polymerization. Facile in Situ Catalytic Synthesis of Silyl-Terminated Polyolefins. *J. Am. Chem. Soc.* **1995**, *117*, 10747–10748.

(13) Brown, H. C.; Khire, U. R.; Narla, G. B-( $\gamma$ -Trimethylsilyl)propargyl)diisopinocampheylborane: A New, Highly Efficient Reagent for the Enantioselective Propargylboration of Aldehydes. Synthesis of Trimethylsilyl-Substituted and Parent  $\alpha$ -Allenic Alcohols in High Optical Purity. *J. Org. Chem.* **1995**, *60* (25), 8130–8131.

(14) Fu, P.-F.; Brard, L.; Li, Y.; Marks, T. J. Regioselection and Enantioselection in Organolanthanide-Catalyzed Olefin Hydrosilylation. A Kinetic and Mechanistic Study. *J. Am. Chem. Soc.* **1995**, *117*, 7157–7168.

(15) Molander, G. A.; Julius, M. Selective hydrosilylation of alkenes catalyzed by an organoyttrium complex. *J. Org. Chem.* **1992**, *57*, 6347–6351.

(16) Sasaki, Y.; Horita, Y.; Zhong, C.; Sawamura, M.; Ito, H. Copper(I)-Catalyzed Regioselective Monoborylation of 1,3-Enynes with an Internal Triple Bond: Selective Synthesis of 1,3-Dienylboronates and 3-Alkynylboronates. *Angew. Chem., Int. Ed.* **2011**, *50*, 2778–2782.

(17) (a) Hajela, S.; Schaefer, W. P.; Bercaw, J. E. Highly electron deficient group 3 organometallic complexes based on the 1,4,7-trimethyl-1,4,7-triazacyclononane ligand system. *J. Organomet. Chem.* **1997**, *532*, 45–53. (b) Ihara, E.; Tanaka, M.; Yasuda, H.; Kanehisa, N.; Maruo, T.; Kai, Y. Coordination of allenyl/propargyl group to samarium(III): the first crystal structures of  $\eta^3$ -allenyl and  $\eta^1$ -propargyl lanthanide complexes. *J. Organomet. Chem.* **2000**, *613*, 26–32. (c) Quiroga Norambuena, V. F.; Heeres, A.; Heeres, H. J.; Meetsma, A.; Teuben, J. H.; Hessen, B. Synthesis, Structure, and Reactivity of Rare-Earth Metallocene  $\eta^3$ -Propargyl/Allenyl Complexes. *Organometallics* **2008**, *27*, 5672–5683. (d) Kaneko, H.; Nagae, H.; Tsurugi, H.; Mashima, K. End-Functionalized Polymerization of 2-Vinylpyridine through Initial C–H Bond Activation of N-Heteroaromatics and Internal Alkynes by Yttrium Ene–Diamido Complexes. *J. Am. Chem. Soc.* **2011**, *133*, 19626–19629. (e) Kefalidis, C. E.; Perrin, L.; Burns, C. J.; Berg, D. J.; Maron, L.; Andersen, R. A. Can a pentamethylcyclopentadienyl ligand act as a proton-relay in f-element chemistry? Insights from a joint experimental/theoretical study. *Dalton Trans.* **2015**, *44*, 2575–2587. (f) Nagae, H.; Kundu, A.; Tsurugi, H.; Mashima, K. Propargylic C(sp<sup>3</sup>)–H Bond Activation for Preparing  $\eta^3$ -Propargyl/Allenyl Complexes of Yttrium. *Organometallics* **2017**, *36*, 3061–3067.