

Strong and Confined Acids Catalyze Asymmetric Intramolecular Hydroarylations of Unactivated Olefins with Indoles

Pinglu Zhang, Nobuya Tsuji, Jie Ouyang, and Benjamin List*



Cite This: *J. Am. Chem. Soc.* 2021, 143, 675–680



Read Online

ACCESS |



Metrics & More



Article Recommendations




Supporting Information


ABSTRACT: In recent years, several organocatalytic asymmetric hydroarylations of activated, electron-poor olefins with activated, electron-rich arenes have been described. In contrast, only a few approaches that can handle *unactivated*, electronically neutral olefins have been reported and invariably require transition metal catalysts. Here we show how an efficient and highly enantioselective catalytic asymmetric intramolecular hydroarylation of aliphatic and aromatic olefins with indoles can be realized using strong and confined IDPi Brønsted acid catalysts. This unprecedented transformation is enabled by tertiary carbocation formation and establishes quaternary stereogenic centers in excellent enantioselectivity and with a broad substrate scope that includes an aliphatic iodide, an azide, and an alkyl boronate, which can be further elaborated into bioactive molecules.

A particularly elegant and atom-economical approach to the catalytic formation of benzylic quaternary stereogenic centers would be via the protonation of a 1,1-disubstituted olefin with a chiral acid, followed by an asymmetric counteranion-directed Friedel–Crafts arylation of the resulting tertiary carbocation (eq 1). However, to the best of our

Design: A perfectly atom-economical catalytic asymmetric hydroarylation of olefins via a tertiary carbocation intermediate



Realization here: Strong and confined acid-catalyzed intramolecular hydroarylation of unactivated olefins



knowledge, such a transformation has not yet been realized.¹ This discrepancy is even more remarkable, given that nonasymmetric, catalytic hydroarylations of unactivated olefins with arenes are common carbon–carbon σ bond-forming reactions and are applied on a multimillion-ton scale.² Previous organocatalytic asymmetric hydroarylations invariably required electronically activated olefins and arenes,³ while transition-metal-catalyzed variants typically relied on covalently bound directing groups.^{4–8} Notable advances toward less biased substrates in intramolecular asymmetric hydroarylations with indoles using π -Lewis acidic Pt- or Ni-catalysts have been reported by Widenhoefer, Peters, and Cramer.^{9–11} Additionally, Hartwig and Meek developed enantioselective Ir- and Rh-catalyzed intermolecular hydroarylations using norbornene and 1-arylbutadienes as olefin components, respectively.^{12,13}


However, the catalytic construction of quaternary stereogenic centers via hydroarylation of unactivated olefins has remained elusive to date, using either transition metals or organic catalysts.^{8b,14} We have recently discovered that strong and confined imidodiphosphorimidate (IDPi) Brønsted acid catalysts can protonate unactivated olefins to engage in highly enantioselective Markovnikov hydroalkoxylations.^{15,16} Our findings suggested the applicability of this unique activation mode for additional transformations that proceed via carbocation intermediates,¹⁷ and an asymmetric hydroarylation of simple olefins became particularly intriguing to us. We now report an intramolecular hydroarylation of unbiased olefins with indoles, furnishing highly enantioenriched tetrahydrocarbazoles possessing a benzylic quaternary stereogenic center (eq 2).

At the onset of our studies, we subjected 1,1-disubstituted olefin **1a** to 2 mol % of various chiral Brønsted acid catalysts at 80 °C for 20 h (Table 1). While weaker acids such as phosphoric acid (CPA) **3** ($pK_a = 12.7$ in MeCN),¹⁸ disulfonimide **4** ($pK_a > 8.4$),^{15b,19,20} imidodiphosphoric acid (IDP) **5** (pK_a ca. 11.5),^{15b,21} iminoimidodiphosphate (*i*IDP) **6** (pK_a ca. 9),^{15b,22} gave poor or moderate conversions, yields, and enantioselectivities (entries 1–4), IDPi catalyst **7a** ($pK_a = 4.5$),²⁰ in contrast, furnished an almost racemic product in 90% yield, suggesting that the key requirement for reactivity is high acidity (entry 5). Installation of biphenyl substituents at the 3,3'-positions of the BINOL backbone, combined with a perfluoronaphthalene-2-sulfonyl “inner core” (IDPi **7c**), quickly revealed a significant increase in enantioselectivity

Received: November 17, 2020

Published: January 5, 2021



Table 1. Reaction Development^a

entry	catalyst	T (°C)	conv. (%)	yield 2a (%)	isom. (%)	er
1	3	80	41	34	5	54:46
2	4	80	47	30	trace	44:56
3	5	80	13	<10	trace	53:47
4	6	80	55	45	9	47:53
5	7a	80	95	90	trace	49:51
6	7b	80	full	93	—	56:44
7	7c	80	full	85	—	88:12
8	7d	80	full	93	—	88:12
9	7e	80	full	93	—	90:10
10 ^b	7e	60	full	95	—	95:5

^aReactions were performed with substrate **1a** (0.02 mmol), catalyst (2 mol %) in methylcyclohexane (CyMe, 0.2 mL); conversions (conv.), yields of **2a**, and olefin isomerizations (isom.) were determined by ¹H NMR analysis with mesitylene as an internal standard; enantiomeric ratios (er's) were measured by HPLC. When the reaction was not carried out in the dark, byproducts and lower yields were observed.
^b48 h.

(entry 7, 88:12 er). Closer inspection of the crystal structure of catalyst **7c** (see Supporting Information) revealed extensive π - π -interactions between the 3,3'-substituents and the aromatic inner core, significantly influencing the shape of the confined, chiral microenvironment. We reasoned that more electron-rich catalyst substituents could potentially stabilize both the π - π -interactions and the critical cationic intermediate. Indeed, with methoxy-substituted catalyst **7d**, yields could be increased (entry 8, 93%). Finally, we identified IDPi **7e** as the optimal catalyst, which, in addition to the 4'-methoxy group, features two *n*-hexyl chains in the 3'- and 5'-positions of the biphenyl system, affording the tetrahydrocarbazole **2a** in both excellent yield and enantioselectivity (entry 10, 95%, 95:5 er).


The outstanding performance of catalyst **7e** was then further challenged with several different substrates to explore the scope of the hydroarylation. In general, the reaction tolerates an array of substituents with differing steric and electronic properties, in addition to sensitive functional groups. As displayed in Table 2, irrespective of the alkyl groups attached to the olefins, the corresponding tetrahydrocarbazoles **2a**–**2g** were formed in excellent yields and enantioselectivities. The

reaction is not limited to aliphatic olefins, but also the phenyl-substituted olefin **1h** gave the corresponding product **2h** in 88% yield and 95:5 er. Methyl substituents at any of the benzo-positions (4, 5, 6, and 7) of the indole core did not hinder the recognition with the catalyst, and products **2i**–**2l** formed with equally high enantioselectivities. The annulated substrates, benzo[*g*]indole and tetrahydrocyclopenta[*g*]indole, also furnished the desired products **2m** and **2n** in excellent yields and enantioselectivities. Furthermore, halogen substituents as potential synthetic handles for future modifications, as well as electron-rich alkoxy substituents, were well tolerated (**2o**–**2s**). Notably, terminal dioxoborolane-, azido-, and iodo-substitution at the alkyl chain of the olefin fragment gave direct and highly enantioselective access to tetrahydrocarbazoles **2t**–**2v**, illustrating the somewhat unexpected functional group tolerance of our strong acid catalysts. The absolute configuration of boronate product **2t** was determined by single-crystal X-ray diffraction, and the configuration of all other products was assigned by analogy.

To further demonstrate the synthetic utility of the hydroarylation products, boronate **2t** was oxidatively hydrolyzed to the corresponding primary alcohol **2w** with complete retention of enantiopurity. Additionally, a C(sp³)-C(sp²) Suzuki–Miyaura cross-coupling with 2-bromonaphthalene provided compound **2x** in 77% yield and, once again, without any erosion of enantiopurity. The subsection of azido-functionalized tetrahydrocarbazole **2u** to a sequential *N*-ethylation, azide reduction, and *N,N*-dimethylation provided the first enantioselective synthesis of ammonium salt **8**, in 41% overall yield and with retention of enantiopurity, the racemic mixture of which is being investigated as an antidepressant.²³

Keen on understanding the underlying mechanism of our hydroarylation, we performed several control experiments (Scheme 1). Moderately acidic chiral Brønsted acid catalysts such as CPAs have been shown to typically operate via a bifunctional, double-hydrogen bonding mechanism.^{18b,c} As a consequence, the enantioselectivity and reactivity of CPA-catalyzed reactions involving a nucleophilic indole can massively erode upon *N*-substitution due to the lack of a recognition and activation element.²⁴ In contrast, we found that *N*-methylated indole **9** readily converted to the corresponding product under our standard conditions, albeit with a significantly reduced enantioselectivity of 71:29 er (Scheme 1A). This observation is consistent with a bifunctional enantiodiscrimination mechanism involving hydrogen bonding interactions in a putative ion-pairing scenario.

Electrophilic aromatic substitutions of indoles have been shown to operate through a direct reaction at C-2 or the more nucleophilic C-3 position, followed by a migratory opening of a spiroindolenine intermediate.^{25,26} We approached the important mechanistic question of how the cyclization occurs experimentally in three different ways (Scheme 1B, C, and D). First, removal of one methylene group from the alkyl tether between indole and olefin in substrate **11** would lead to the corresponding tetrahydrocyclopenta[*b*]indole **12**, the formation of which, if nucleophilic attack would proceed directly from C-2, would be expected to be kinetically at least as favorable as the corresponding six-membered ring formation.²⁷ However, even after stirring olefin **11** at 120 °C for 2 d in the presence of 5 mol % of catalyst **7e**, product **12** could not be observed, excluding direct reaction at C-2. This result is consistent with the initial C-3 spirocyclization mechanism, which in this specific case is not viable because

Table 2. Scope of the Reaction^a

^aReactions were carried out with 0.1–0.2 mmol of substrates **1**, catalyst **7e** (2 mol %) in cyclohexane (0.1 M) at 60 °C unless otherwise noted. Yields are for the isolated compounds. The enantiomeric ratios (er) were determined by HPLC analysis. ^bReaction was run at 60 °C for 3 d. ^cReaction was run with 3 mol % catalyst at 65 °C for 3 d. ^dNaBO₃·H₂O (5 equiv), THF/H₂O 1:1, rt, 2 h, 92% yield. ^e2-Bromonaphthalene (1.05 equiv), Pd(OAc)₂ (2.5 mol %), RuPhos (5 mol %), 85 °C, 24 h, 77% yield. ^f(i) NaH (1.2 equiv), iodoethane (1.2 equiv), 0 °C to rt, DMF, 2 h, 85% yield; (ii) NaBH₄ (2 equiv), CoCl₂·6H₂O (10 mol %), MeOH, 0 °C, 30 min, 84% yield; (iii) NaBH₃CN (2 equiv), MeOH, HCHO (37 wt % in H₂O), rt, 3 h, 58% yield; (iv) HCl (1 M in Et₂O), quant. THF = tetrahydrofuran; DMF = dimethylformamide.


of the high ring strain of the required four-membered spirocyclic intermediate. Second, addition of a methylene group to the linkage in substrate **13** afforded the corresponding seven-membered hexahydrocyclohepta[*b*]indole **14** in moderate yield and 76:24 er, consistent with a six-membered spirocyclic intermediate. Third, we subjected terminal olefin **15** to our reaction conditions using IDPi catalysts **7a** and **7e**, which would undergo cyclization via a secondary carbocation intermediate. Remarkably, both reactions went to full conversion, although catalyst **7a** required a slightly elevated temperature to reach completion. Even more interestingly, IDPi **7a** provided the tetrahydrocarbazole product as a regioisomeric mixture of **16a/16b** = 6.7/1,²⁸ strongly suggesting an initial spirocyclization followed by competitive migration of either methylene or methide fragment. Unfortunately, ¹H NMR investigations of the cyclization of substrate **1a** under standard conditions (see Supporting Information) did not lead to the detection of a spiroindolenine intermediate, presumably due to a relatively fast migratory process restoring indole aromaticity.

Based on these experiments, we propose a catalytic cycle as depicted in Scheme 1E. Accordingly, the reaction is initiated

by a protonation of the olefin to give ion-pair I, followed by a nucleophilic attack of the indole directed by the chiral counteranion, which is coordinating to the substrate via hydrogen bonding to the indole N–H moiety. This spirocyclization gives rise to the spiroindolenine cation in association with the IDPi anion (ion-pair II). Rapid suprafacial and stereoretentive migration of the most electron-rich alkyl fragment then liberates tetrahydrocarbazole **2** and, upon proton transfer, restores catalyst **7e**.^{25b}

We speculate that the remarkable substrate scope and functional group tolerance of our system could result from a confinement effect, according to which the catalyst recognizes (or “binds”) the transition state of the cyclization. In contrast, the olefin substituent is mostly positioned outside of the chiral, confined active site, well within the solvent. The transition-state-binding scenario is also typical for even more confined enzymes, with the important difference that our catalyst’s incomplete confinement enables a significant “promiscuity”. Conceptually, these features position our confined acid catalysts in-between small-molecule catalysts on the one end and macromolecular catalysts with a substrate-specific active site on the other end of the spectrum. The peculiar behavior of

Scheme 1. Mechanistic Study



^aReaction was carried out with 0.2 mmol of **9**, catalyst **7e** (2 mol %) in cyclohexane (0.1 M); yield is of for the isolated compound; enantiomeric ratios (er) were measured by HPLC. ^bReactions were performed with substrate **11**, **13**, **15** (0.02 mmol); conversions, yields, and regioisomeric ratios were determined by ¹H NMR analysis with mesitylene as an internal standard. ^c100 °C with 4 mol % **7a** in methylcyclohexane; 60 °C with 2 mol % **7a** in cyclohexane, trace product. ^d60 °C with 2 mol % **7e** in cyclohexane.

our confined catalysts has also been displayed in other reactions, most notably in our recent hydroalkoxylation reaction, which has a similar specificity with regard to the reacting subunit, while tolerating a broad range of (functionalized) substituents.¹⁶

We have developed a perfectly atom-economical and highly enantioselective catalytic intramolecular hydroarylation of unactivated olefins with indoles. Our reaction is enabled via olefin protonation with highly acidic and confined chiral Brønsted acid catalysts and provides access to valuable enantiopure tetrahydrocarbazoles.²⁹ The previously unattainable activation of 1,1-disubstituted olefin via protonation to the corresponding tertiary carbocation suggests obvious and significant potential for other hydrofunctionalization reactions that provide quaternary stereogenic center-bearing products and related compounds.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.0c12042>.

Crystallographic data for compound **2t** (CIF)

Crystallographic data for compound **7c** (CIF)

Experimental details and analytical data for all new compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Benjamin List – Max-Planck-Institut für Kohlenforschung, 45470 Mülheim an der Ruhr, Germany; Institute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo 001-0021, Japan; orcid.org/0000-0002-9804-599X; Email: list@kofo.mpg.de

Authors

Pinglu Zhang – Max-Planck-Institut für Kohlenforschung, 45470 Mülheim an der Ruhr, Germany

Nobuya Tsuji – Max-Planck-Institut für Kohlenforschung, 45470 Mülheim an der Ruhr, Germany; Institute for Chemical Reaction Design and Discovery (WPI-ICReDD), Hokkaido University, Sapporo 001-0021, Japan

Jie Ouyang – Max-Planck-Institut für Kohlenforschung, 45470 Mülheim an der Ruhr, Germany

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacs.0c12042>

Notes

The authors declare no competing financial interest. Crystallographic data for compounds **2t** (CCDC 2044898) and **7c** (CCDC 2044899) can be downloaded free of charge from the Cambridge Crystallographic Data Centre (www.ccdc.cam.ac.uk).

■ ACKNOWLEDGMENTS

Generous support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Leibniz Award to B.L. and under Germany's Excellence Strategy–EXC 2033–390677874–RESOLV, and the European Research Council (ERC, European Union's Horizon 2020 research and innovation program “C–H Acids for Organic Synthesis, CHAOS” Advanced Grant Agreement No. 694228) and the Alexander von Humboldt Foundation for a fellowship to P.Z. is gratefully acknowledged. We further thank Dr. Markus Leutzsch for NMR studies, Benjamin Mitschke for his help during the preparation of this manuscript, Lukas Münzer for his contributions in initial studies, and several members of the group for crowd reviewing this manuscript. We also thank the

technicians of our group and the members of our NMR, MS, X-ray, and chromatography groups for their excellent service.

REFERENCES

- (1) Properzi, R.; Kaib, P. S. J.; Leutzsch, M.; Pupo, G.; Mitra, R.; De, C. K.; Song, L.; Schreiner, P. R.; List, B. Catalytic enantiocontrol over a non-classical carbocation. *Nat. Chem.* **2020**, *12*, 1174–1179.
- (2) (a) Olah, G. A.; Molnár, A. *Hydrocarbon Chemistry*; Wiley-Interscience: New York, 2003. (b) Perego, C.; Ingallina, P. Combining alkylation and transalkylation for alkylaromatic production. *Green Chem.* **2004**, *6*, 274–279. (c) Andreatta, J. R.; McKeown, B. A.; Gunnoe, T. B. Transition metal catalyzed hydroarylation of olefins using unactivated substrates: Recent developments and challenges. *J. Organomet. Chem.* **2011**, *696*, 305–315. (d) Cai, X.; Tohti, A.; Ramirez, C.; Harb, H.; Fettinger, J. C.; Hratchian, H. P.; Stokes, B. J. Dispersion-Controlled Regioselective Acid-Catalyzed Intramolecular Hydroindolization of *cis*-Methindolylstyrenes To Access Tetrahydrobenzo[*cd*]indoles. *Org. Lett.* **2019**, *21*, 1574–1577.
- (3) (a) Paras, N. A.; MacMillan, D. W. C. New Strategies in Organic Catalysis: The First Enantioselective Organocatalytic Friedel–Crafts Alkylation. *J. Am. Chem. Soc.* **2001**, *123*, 4370–4371. (b) You, S.-L.; Cai, Q.; Zeng, M. Chiral Brønsted acid catalyzed Friedel–Crafts alkylation reactions. *Chem. Soc. Rev.* **2009**, *38*, 2190–2201. (c) Terrasson, V.; Marcia de Figueiredo, R.; Campagne, J. M. Organocatalyzed Asymmetric Friedel–Crafts Reactions. *Eur. J. Org. Chem.* **2010**, *2010*, 2635–2655. (d) Correia, J. T. M.; List, B.; Coelho, F. Catalytic Asymmetric Conjugate Addition of Indolizines to α,β -Unsaturated Ketones. *Angew. Chem., Int. Ed.* **2017**, *56*, 7967–7970.
- (4) For reviews, see: (a) Dong, Z.; Ren, Z.; Thompson, S. J.; Xu, Y.; Dong, G. Transition-Metal-Catalyzed C–H Alkylation Using Alkenes. *Chem. Rev.* **2017**, *117*, 9333–9403. (b) Coombs, J. R.; Morken, J. P. Catalytic Enantioselective Functionalization of Unactivated Terminal Alkenes. *Angew. Chem., Int. Ed.* **2016**, *55*, 2636–2649. (c) Newton, C. G.; Wang, S.-G.; Oliveira, C. C.; Cramer, N. Catalytic Enantioselective Transformations Involving C–H Bond Cleavage by Transition-Metal Complexes. *Chem. Rev.* **2017**, *117*, 8908–8976.
- (5) For selected examples using a directing group in Ir-catalysis, see: (a) Aufdenblatten, R.; Diezi, S.; Togni, A. Iridium(I)-Catalyzed Asymmetric Intermolecular Hydroarylation of Norbornene with Benzamide. *Monatsh. Chem.* **2000**, *131*, 1345–1350. (b) Shirai, T.; Yamamoto, Y. Cationic Iridium/S-Me-BIPAM-Catalyzed Direct Asymmetric Intermolecular Hydroarylation of Bicycloalkenes. *Angew. Chem., Int. Ed.* **2015**, *54*, 9894–9897. (c) Grélaud, S.; Cooper, P.; Feron, L. J.; Bower, J. F. Branch-Selective and Enantioselective Iridium-Catalyzed Alkene Hydroarylation via Anilide-Directed C–H Oxidative Addition. *J. Am. Chem. Soc.* **2018**, *140*, 9351–9356. (d) Shibata, T.; Ryu, N.; Takano, H. Iridium-Catalyzed Intramolecular Enantioselective C–H Alkylation at the C-2 Position of *N*-Alkenylindoles. *Adv. Synth. Catal.* **2015**, *357*, 1131–1135.
- (6) For selected examples using directing group in Rh-catalysis, see: (a) Thalji, R. K.; Ellman, J. A.; Bergman, R. G. Highly Efficient and Enantioselective Cyclization of Aromatic Imines via Directed C–H Bond Activation. *J. Am. Chem. Soc.* **2004**, *126*, 7192–7193. (b) Harada, H.; Thalji, R. K.; Bergman, R. G.; Ellman, J. A. Enantioselective Intramolecular Hydroarylation of Alkenes via Directed C–H Bond Activation. *J. Org. Chem.* **2008**, *73*, 6772–6779. (c) Ye, B.; Cramer, N. Chiral Cyclopentadienyl Ligands as Stereocontrolling Element in Asymmetric C–H Functionalization. *Science* **2012**, *338*, 504–506.
- (7) For selected examples using directing group in Co-catalysis, see (a) Lee, P.-S.; Yoshikai, N. Cobalt-Catalyzed Enantioselective Directed C–H Alkylation of Indole with Styrenes. *Org. Lett.* **2015**, *17*, 22–25. (b) Pesciaoli, F.; Dhawa, U.; Oliveira, J. C. A.; Yin, R.; John, M.; Ackermann, L. Enantioselective Cobalt(III)-Catalyzed C–H Activation Enabled by Chiral Carboxylic Acid Cooperation. *Angew. Chem., Int. Ed.* **2018**, *57*, 15425–15429.
- (8) For selected examples Using directing group in other metal catalysis, see: (a) Loup, J.; Zell, D.; Oliveira, J. C. A.; Keil, H.; Stalke, D.; Ackermann, L. Asymmetric Iron-Catalyzed C–H Alkylation Enabled by Remote Ligand meta-Substitution. *Angew. Chem., Int. Ed.* **2017**, *56*, 14197–14201. (b) Li, G.; Liu, Q.; Vasamsetty, L.; Guo, W.; Wang, J. Ruthenium(II)-Catalyzed Asymmetric Inert C–H Bond Activation Assisted by a Chiral Transient Directing Group. *Angew. Chem., Int. Ed.* **2020**, *59*, 3475–3479. (c) Song, G.; O, W. W. N.; Hou, Z. Enantioselective C–H Bond Addition of Pyridines to Alkenes Catalyzed by Chiral Half-Sandwich Rare-Earth Complexes. *J. Am. Chem. Soc.* **2014**, *136*, 12209–12212.
- (9) (a) Liu, C.; Han, X.; Wang, X.; Widenhoefer, R. A. Platinum-Catalyzed Intramolecular Alkylation of Indoles with Unactivated Olefins. *J. Am. Chem. Soc.* **2004**, *126*, 3700–3701. (b) Han, X.; Widenhoefer, R. A. Platinum-Catalyzed Intramolecular Asymmetric Hydroarylation of Unactivated Alkenes with Indoles. *Org. Lett.* **2006**, *8*, 3801–3804.
- (10) Huang, H.; Peters, R. A. Highly Strained Planar-Chiral Platinum Cycle for Catalytic Activation of Internal Olefins in the Friedel–Crafts Alkylation of Indoles. *Angew. Chem., Int. Ed.* **2009**, *48*, 604–606.
- (11) Diesel, J.; Grosheva, D.; Kodama, S.; Cramer, N. A Bulky Chiral *N*-Heterocyclic Carbene Nickel Catalyst Enables Enantioselective C–H Functionalizations of Indoles and Pyrroles. *Angew. Chem., Int. Ed.* **2019**, *58*, 11044–11048.
- (12) Sevov, C. S.; Hartwig, J. F. Iridium-Catalyzed Intermolecular Asymmetric Hydroheteroarylation of Bicycloalkenes. *J. Am. Chem. Soc.* **2013**, *135*, 2116–2119.
- (13) Marcum, J. S.; Roberts, C. C.; Manan, R. S.; Cervarich, T. N.; Meek, S. J. Chiral Pincer Carbodicarbene Ligands for Enantioselective Rhodium-Catalyzed Hydroarylation of Terminal and Internal 1,3-Dienes with Indoles. *J. Am. Chem. Soc.* **2017**, *139*, 15580–15583.
- (14) (a) Lou, S.-J.; Mo, Z.; Nishiura, M.; Hou, Z. Construction of All-Carbon Quaternary Stereocenters by Scandium-Catalyzed Intramolecular C–H Alkylation of Imidazoles with 1,1-Disubstituted Alkenes. *J. Am. Chem. Soc.* **2020**, *142*, 1200–1205. (b) Shinde, V. S.; Mane, M. V.; Cavallo, L.; Rueping, M. Iridium-Catalyzed Enantioselective Hydroarylation of Alkenes through C–H bond Activation: Experiment and Computation. *Chem. - Eur. J.* **2020**, *26*, 8308–8313. (c) Ye, B.; Donets, P. A.; Cramer, N. Chiral Cp-Rhodium(III)-Catalyzed Asymmetric Hydroarylations of 1,1-Disubstituted Alkenes. *Angew. Chem., Int. Ed.* **2014**, *53*, 507–511.
- (15) (a) Kaib, P. S. J.; Schreyer, L.; Lee, S.; Properzi, R.; List, B. Extremely Active Organocatalysts Enable a Highly Enantioselective Addition of Allyltrimethylsilane to Aldehydes. *Angew. Chem., Int. Ed.* **2016**, *55*, 13200–13203. (b) Schreyer, L.; Properzi, R.; List, B. IDPi Catalysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 12761–12777.
- (16) Tsuji, N.; Kennemur, J. L.; Buyck, T.; Lee, S.; Prévost, S.; Kaib, P. S. J.; Bykov, D.; Farès, C.; List, B. Activation of olefins via asymmetric Brønsted acid catalysis. *Science* **2018**, *359*, 1501–1505.
- (17) For examples of substitution reactions proceeding via tertiary carbocations, see: (a) Isomura, M.; Petrone, D. A.; Carreira, E. M. Coordination-Induced Stereocontrol over Carbocations: Asymmetric Reductive Deoxygenation of Racemic Tertiary Alcohols. *J. Am. Chem. Soc.* **2019**, *141*, 4738–4748. (b) Braun, M.; Kotter, W. Titanium(IV)-Catalyzed Dynamic Kinetic Asymmetric Transformation of Alcohols, Silyl Ethers, and Acetals under Carbon Allylation. *Angew. Chem., Int. Ed.* **2004**, *43*, 514–517. (c) Zhao, W.; Wang, Z.; Chu, B.; Sun, J. Enantioselective Formation of All-Carbon Quaternary Stereocenters from Indoles and Tertiary Alcohols Bearing A Directing Group. *Angew. Chem., Int. Ed.* **2015**, *54*, 1910–1913. (d) Wendlandt, A. E.; Vangal, P.; Jacobsen, E. N. Quaternary Stereocenters via an Enantioconvergent Catalytic S_N1 Reaction. *Nature* **2018**, *556*, 447–451. Also see: (e) Kikuchi, J.; Terada, M. Enantioselective Addition Reaction of Azlactones with Styrene Derivatives Catalyzed by Strong Chiral Brønsted Acids. *Angew. Chem., Int. Ed.* **2019**, *58*, 8458–8462.
- (18) (a) Kaupmees, K.; Tolstoluzhsky, N.; Raja, S.; Rueping, M.; Leito, I. On the Acidity and Reactivity of Highly Effective Chiral Brønsted Acid Catalysts: Establishment of an Acidity Scale. *Angew. Chem., Int. Ed.* **2013**, *52*, 11569–11572. (b) Akiyama, T. Stronger Brønsted acids. *Chem. Rev.* **2007**, *107*, 5744–5758. (c) Akiyama, T.;

Mori, K. Stronger Brønsted acids: Recent Progress. *Chem. Rev.* **2015**, *115*, 9277–9306.

(19) (a) García-García, P.; Lay, F.; García-García, P.; Rabalakos, C.; List, B. A Powerful Chiral Counteranion Motif for Asymmetric Catalysis. *Angew. Chem., Int. Ed.* **2009**, *48*, 4363–4366. (b) James, T.; van Gemmeren, M.; List, B. Development and Applications of Disulfonimides in Enantioselective Organocatalysis. *Chem. Rev.* **2015**, *115*, 9388–9409. (c) Wakchaure, V. N.; Obradors, C.; List, B. Chiral Brønsted Acids Catalyze Asymmetric Additions to Substrates that Are Already Protonated: Highly Enantioselective Disulfonimide-Catalyzed Hantzsch Ester Reductions of NH–Imine Hydrochloride Salts. *Synlett* **2020**, *31*, 1707–1712. (d) Zhang, Z.; Klusmann, M.; List, B. Kinetic Study of Disulfonimide-Catalyzed Cyanosilylation of Aldehydes by Using a Method of Progress Rates. *Synlett* **2020**, *31*, 1593–1597.

(20) Bae, H. Y.; Höfler, D.; Kaib, P. S. J.; Kasaplar, P.; De, C. K.; Döhning, A.; Lee, S.; Kaupmees, K.; Leito, I.; List, B. Approaching sub-ppm-level asymmetric organocatalysis of a highly challenging and scalable carbon–carbon bond forming reaction. *Nat. Chem.* **2018**, *10*, 888–894.

(21) Corić, I.; List, B. Asymmetric spiroacetalization catalysed by confined Brønsted acids. *Nature* **2012**, *483*, 315–319.

(22) Liu, L.; Kaib, P. S. J.; Tap, A.; List, B. A General Catalytic Asymmetric Prins Cyclization. *J. Am. Chem. Soc.* **2016**, *138*, 10822–10825.

(23) (a) Pugsley, T. A.; Lippmann, W. Effects of 1-alkyl-1,2,3,4-tetrahydrocarbazole-1-ethanamines and related compounds, potential antidepressants, on biogenic amine uptake mechanisms. *Experientia* **1977**, *33*, 57–59. (b) Asselin, A. A.; Humber, L. G.; Komlossy, J.; Charest, M. P. Cycloalkanoindoles. 2. 1-Alkyl-1,2,3,4-tetrahydrocarbazole-1-ethanamines and related compounds. Potential antidepressants. *J. Med. Chem.* **1976**, *19*, 792–797.

(24) (a) Sheng, Y.-F.; Gu, Q.; Zhang, A.-J.; You, S.-L. Chiral Brønsted Acid-Catalyzed Asymmetric Friedel–Crafts Alkylation of Pyrroles with Nitroolefins. *J. Org. Chem.* **2009**, *74*, 6899–6901. (b) Itoh, J.; Fuchibe, K.; Akiyama, T. Chiral Phosphoric Acid Catalyzed Enantioselective Friedel–Crafts Alkylation of Indoles with Nitroalkenes: Cooperative Effect of 3 Å Molecular Sieves. *Angew. Chem., Int. Ed.* **2008**, *47*, 4016–4018. (c) Mori, K.; Wakazawa, M.; Akiyama, T. Stereoselective construction of all-carbon quaternary center by means of chiral phosphoric acid: highly enantioselective Friedel–Crafts reaction of indoles with β,β -disubstituted nitroalkenes. *Chem. Sci.* **2014**, *5*, 1799–1803.

(25) (a) Wu, Q.-F.; Zheng, C.; You, S.-L. Enantioselective Synthesis of Spiro Cyclopentane-1,3'-indoles and 2,3,4,9-Tetrahydro-1H-carbazoles by Iridium-Catalyzed Allylic Dearomatization and Stereospecific Migration. *Angew. Chem., Int. Ed.* **2012**, *51*, 1680–1683. (b) Zheng, C.; Wu, Q.-F.; You, S.-L. A Combined Theoretical and Experimental Investigation into the Highly Stereoselective Migration of Spiroindolenines. *J. Org. Chem.* **2013**, *78*, 4357–4365. (c) Zheng, C.; You, S.-L. Exploring the Chemistry of Spiroindolenines by Mechanistically-Driven Reaction Development: Asymmetric Pictet–Spengler-type Reactions and Beyond. *Acc. Chem. Res.* **2020**, *53*, 974–987.

(26) (a) Jackson, A. H.; Naidoo, B.; Smith, P. Electrophilic substitution in indoles—IV: The cyclization of indolylbutanol to tetrahydrocarbazole. *Tetrahedron* **1968**, *24*, 6119–6129. (b) Casnati, G.; Dossena, A.; Pochini, A. Electrophilic substitution in indoles: direct attack at the 2-position of 3-alkylindoles. *Tetrahedron Lett.* **1972**, *13*, 5277–5280.

(27) (a) Baldwin, J. E. Rules for ring closure. *J. Chem. Soc., Chem. Commun.* **1976**, 734–736. (b) Casadei, M. A.; Galli, C.; Mandolini, L. Ring-closure reactions. 22. Kinetics of cyclization of diethyl (ω -Bromoalkyl)malonates in the range of 4- to 21-membered rings. Role of ring strain. *J. Am. Chem. Soc.* **1984**, *106*, 1051–1056. (c) Alabugin, I. V.; Gilmore, K. Finding the right path: Baldwin “Rules for Ring Closure” and stereoelectronic control of cyclizations. *Chem. Commun.* **2013**, *49*, 11246–11250.

(28) The configuration of **16a** was assigned by comparing specific rotation of analogy in ref **25a**. **16a** 69:31 e.r.; $[\alpha]_{\text{D}}^{25} = 8.0$ (c 0.1, CHCl_3).

(29) (a) Kochanowska-Karamyan, A. J.; Hamann, M. T. Marine Indole Alkaloids: Potential New Drug Leads for the Control of Depression and Anxiety. *Chem. Rev.* **2010**, *110*, 4489–4497. (b) Tan, F.; Cheng, H.-G. Catalytic asymmetric synthesis of tetrahydrocarbazoles. *Chem. Commun.* **2019**, *55*, 6151–6164.